






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 487/04, A61K 31/495 // (C07D 487/04, 241:00, 209:00)	A1	(11) International Publication Number: WO 97/23482 (43) International Publication Date: 3 July 1997 (03.07.97)
(21) International Application Number: PCT/IB96/01192 (22) International Filing Date: 6 November 1996 (06.11.96) (30) Priority Data: 60/008,988 21 December 1995 (21.12.95) US (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): SANNER, Mark, A. [US/US]; 8 Hillcrest Drive, Old Saybrook, CT 06475 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		(81) Designated States: AU, BG, BR, BY, CA, CN, CZ, HU, IL, IS, JP, KR, KZ, LK, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: 2,7-SUBSTITUTED OCTAHYDRO-PYRROLO[1,2-A]PYRAZINE DERIVATIVES**(57) Abstract**

This invention relates to compounds of formula (I), wherein R₁ is phenyl, naphthyl, benzoxazolonyl, indolyl, indolonyl, benzimidazolyl, quinolyl, furyl, benzofuryl, thienyl, benzothienyl, oxazolyl, benzoxazolyl; R₂ is H or (C₁-C₆)alkyl; R₃ is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, or pyridazinyl; X is O, S, SO, SO₂, NR₄, C=O, CH(OH), CHR₄, (a), (b), (c), or (d); m is 0, 1 or 2; n is 0, 1 or 2. The compounds of this invention are ligands for dopamine receptor subtypes, especially the dopamine D₄ receptor, and are therefore useful in the treatment of disorders of the dopamine system.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

5

2,7-SUBSTITUTED OCTAHYDRO-
PYRROLO[1,2-A]PYRAZINE DERIVATIVES

The present invention relates to novel pharmacologically active 2,7-substituted octahydro-1H-pyrrolo[1,2-a]pyrazine derivatives, their acid addition salts, and certain precursors thereto. The compounds of this invention are ligands for dopamine receptor subtypes, especially the dopamine D₄ receptor, and are therefore useful in the treatment of disorders of the dopamine system.

Background of the Invention

Molecular biological techniques have revealed the existence of several subtypes of the dopamine receptor. The dopamine D₁ receptor subtype has been shown to occur in the least two discrete forms. Two forms of the D₂ receptor subtype, and at least one form of the D₃ receptor subtype, have also been discovered. More recently, the D₄ (Van Tol et al., Nature (London), 1991, 350, 610) and D₅ (Sunahara et al., Nature (London), 1991, 350, 614) receptor subtypes have been described.

The compounds of the present invention, being ligands for dopamine receptor subtypes, especially the dopamine D₄ receptor, are accordingly of use in the treatment or prevention of disorders of the dopamine system.

Dopamine D₄ receptors are more prevalent in the brains of schizophrenic patients (Seeman, et al. Nature, 1993, 365, 441) relative to normal controls. Dopamine receptor antagonists are useful for the treatment of psychotic disorders, such as schizophrenia, and the compounds of the present invention, being ligands for dopamine receptor subtypes, especially the dopamine D₄ receptor, are accordingly of use in the treatment or prevention of psychotic disorders, especially affective psychosis, schizophrenia, and schizoaffective disorders.

Since dopamine receptors control a great number of pharmacological events and, on the other hand, not all these events are presently known, there is a possibility that compounds that act on the dopamine D₄ receptor may exert a wide range of therapeutic effects in animals.

WO 94/10162 (published May 11, 1994) and WO 94/10145 (published May 11, 1994) report that dopamine ligands are of use in the treatment and/or prevention of disorders of the dopamine system, including schizophrenia, nausea, Parkinson's disease, tardive dyskinesias and extrapyramidal side-effects associated with

-2-

treatment by conventional neuroleptic agents, neuroleptic malignant syndrome, and disorders of hypothalamic-pituitary function such as hyperprolactinaemia and amenorrhoea.

Upper gastrointestinal tract motility is believed to be under the control of the dopamine system. The compounds according to the present invention may thus be of use in the prevention and/or treatment of gastrointestinal disorders, and the facilitation of gastric emptying.

Dependence-inducing agents such as cocaine and amphetamine have been shown to interact with the dopamine system. Compounds capable of counteracting this effect, including the compounds in accordance with the present invention, may accordingly be of value in the prevention or reduction of dependence on a dependence-inducing agent.


Dopamine is known to be a peripheral vasodilator; for example, it has been shown to exert a dilatory effect on the renal vascular bed. This implies that the compounds of the present invention may be beneficial in controlling vascular blood flow.

The localization of dopamine receptor mRNA in rat heart and large vessels has been noted. This suggests a role for dopamine receptor ligands in controlling cardiovascular function, either by affecting cardiac and smooth muscle contractility or by modulating the secretion of vasoactive substances. The compounds according to the present invention may therefore be of assistance in the prevention and/or treatment of such conditions as hypertension and congestive heart failure.

The presence of D4 receptor mRNA in mouse retina has been noted (Cohen, et al. *Proc. Nat. Acad. Sci.*, 1992, 89, 12093), suggesting that dopamine and D4 receptors play a role in ocular function. The compounds of this invention may therefore be useful in the treatment of ocular disorders. Furthermore, D4 receptors influence melatonin biosynthesis in chick retina (Zawilska, Nowak, *Neuroscience Lett.*, 1994, 166, 203), and since melatonin has been used for the treatment of sleep disorders, the compounds of this invention may be useful for the treatment of sleep disorders as well.

Diafi, et al. (*J. Het. Chem.*, 1990, 27, 2181) describe certain 2,7-substituted pyrrolo[1,2-a]pyrazine derivatives of compound II for which no biological activity is claimed.


-3-



5

Nakamura, et al. (Jpn. Pat. Appl. JP05097819 A2 930420, published April 20, 1993) describe derivatives and analogs of compound III which are antagonists of platelet activating factor (PAF) for the treatment of asthma and cardiovascular diseases.

10




15

Summary of the Invention

This invention relates to a compound of formula I

20



wherein

25 R_1 is phenyl, naphthyl, benzoxazolonyl, indolyl, indolonyl, benzimidazolyl, quinolyl, furyl, benzofuryl, thienyl, benzothieryl, oxazolyl, benzoxazolyl;

R_2 is H or (C₁-C₆)alkyl;

R_3 is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, or pyridazinyl;

R_4 is H or (C₁-C₆)alkyl;

30


R_5 is H or (C₁-C₆)alkyl;

wherein each group of R_1 and R_3 may be independently and optionally substituted with one to four substituents independently selected from the groups

-4-

consisting of fluoro, chloro, bromo, iodo, cyano, nitro, thiocyano, -SR₄, -SOR₄, -SO₂R₄, -NHSO₂R₄, -(C₁-C₆)alkoxy, -NR₄R₅, -NR₄COR₅, -CONR₄R₅, phenyl, -COR₄, -COOR₄, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl substituted with one to six halogens, -(C₃-C₆)cycloalkyl, and trifluoromethoxy;

5 X is O, S, SO, SO₂, NR₄, C=O, CH(OH), CHR₄,



m is 0, 1 or 2;

n is 0, 1 or 2;

all stereoisomers thereof; or

10 a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₁ is phenyl, naphthyl, benzoxazolonyl, indolyl, indolonyl, benzimidazolyl, or quinolyl;

wherein R₁ and R₃ may be independently substituted with up to three

15 substituents independently selected from the group consisting of fluoro, chloro, bromo, iodo, cyano, -NR₄R₅, -(C₁-C₆)alkoxy, -COOR₄, -CONR₄R₅, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl substituted with one to six halogens, -(C₃-C₆)cycloalkyl, and trifluoromethoxy;

R₂ is H or CH₃;

20 X is O, C=O, CHOH, -C(=O)O-, or CH₂;

m is 0 or 1;

n is 0 or 1; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

25 R₁ is phenyl or substituted phenyl;

R₃ is substituted or unsubstituted phenyl, pyridinyl, or pyrimidinyl;

X is O, -C(=O)O-, or CH₂; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

30 R₂ is H;

X is O;

-5-

m is 0;

n is 1; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

5 R₂ is H;

X is O;

m is 1;

n is 0; or

a pharmaceutically acceptable salt thereof.

10 In another aspect, this invention relates to compounds of formula I wherein

R₂ is H;

X is -C(=O)O-;

m is 0;

n is 0; or

15 a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

R₃ is chloropyridinyl; or

a pharmaceutically acceptable salt thereof.

20 In another aspect, this invention relates to compounds of formula I wherein

R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

R₃ is fluoropyrimidinyl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

25 R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

R₃ is chloropyridinyl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

30 R₃ is fluoropyrimidinyl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

-6-

R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

R₃ is chloropyridinyl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

5 R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

R₃ is fluoropyrimidinyl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₃ is 5-chloro-pyridin-2-yl; or

10 a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₃ is 5-fluoro-pyrimidin-2-yl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

15 R₃ is 5-chloro-pyridin-2-yl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₃ is 5-fluoro-pyrimidin-2-yl; or

a pharmaceutically acceptable salt thereof.

20 In another aspect, this invention relates to compounds of formula I wherein

R₃ is 5-chloro-pyridin-2-yl; or

a pharmaceutically acceptable salt thereof.

In another aspect, this invention relates to compounds of formula I wherein

R₃ is 5-fluoro-pyrimidin-2-yl; or

25 a pharmaceutically acceptable salt thereof.

Preferred compounds of the invention are:

(7S,8aS)-7-(4-fluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;

30 (7S,8aS)-7-(3,5-difluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;

(7S,8aS)-7-(3-cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;

-7-

- (7S,8aS)-7-(4-cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- (7S,8aS)-7-(4-fluorobenzyl)oxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- 5 (7S,8aS)-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate;
- (7S,8aS)-7-(4-fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- (7S,8aS)-7-(3,5-difluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
- 10 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- (7S,8aS)-7-(3-cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
- 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- (7S,8aS)-7-(4-cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
- 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- 15 (7S,8aS)-7-(4-fluorobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine;
- (7S,8aS)-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate;
- (7S,8aS)-7-(3-cyanobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-
- 20 octahydro-pyrrolo[1,2-a]pyrazine;
- and pharmaceutically acceptable salts thereof.

This invention also relates to a method for treating or preventing disorders of the dopamine system in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

This invention also relates to a method for treating or preventing psychotic disorders such as affective psychosis, schizophrenia, and schizoaffective disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

This invention also relates to a method for treating or preventing movement disorders such as extrapyramidal side effects from neuroleptic agents, neuroleptic malignant syndrome, tardive dyskinesia, or Gilles De La Tourette's syndrome in a

-8-

mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

5 This invention also relates to a method for treating or preventing movement disorders such as Parkinson's disease or Huntington's disease in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

10 This invention also relates to a method for treating or preventing gastrointestinal disorders such as gastric acid secretion in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

15 This invention also relates to a method for treating or preventing gastrointestinal disorders such as emesis in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

20 This invention also relates to a method for treating or preventing chemical abuse, chemical dependencies or substance abuse in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

25 This invention also relates to a method for treating or preventing vascular and cardiovascular disorders such as congestive heart failure and hypertension in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

30 This invention also relates to a method for treating or preventing ocular disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

-9-

This invention also relates to a method for treating or preventing sleep disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

5 This invention also relates to a pharmaceutical composition for treating or preventing disorders of the dopamine system in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

10 This invention also relates to a pharmaceutical composition for treating or preventing psychotic disorders such as affective psychosis, schizophrenia, and schizoaffective disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing
15 such disorder.

 This invention also relates to a pharmaceutical composition for treating or preventing movement disorders such as extrapyramidal side effects from neuroleptic agents, neuroleptic malignant syndrome, tardive dyskinesia, or Gilles De La Tourette's syndrome in a mammal, including a human, comprising administering to
20 said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

 This invention also relates to a pharmaceutical composition for treating or preventing movement disorders such as Parkinson's disease or Huntington's disease in a mammal, including a human, comprising administering to said mammal an
25 amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

 This invention also relates to a pharmaceutical composition for treating or preventing gastrointestinal disorders such as gastric acid secretion in a mammal, including a human, comprising administering to said mammal an amount of a
30 compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

 This invention also relates to a pharmaceutical composition for treating or preventing gastrointestinal disorders such as emesis in a mammal, including a

-10-

human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

This invention also relates to a pharmaceutical composition for treating or preventing chemical abuse, chemical dependencies or substance abuse in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.


This invention also relates to a pharmaceutical composition for treating or preventing vascular and cardiovascular disorders such as congestive heart failure and hypertension in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

This invention also relates to a pharmaceutical composition for treating or preventing ocular disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

This invention also relates to a pharmaceutical composition for treating or preventing sleep disorders in a mammal, including a human, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

In another aspect, this invention relates to compounds of formula IV

25



wherein

30

n is 0 or 1;

Y is CH or N;

Z is chloro or fluoro;

all stereoisomers thereof;

-11-

which are useful intermediates for the preparation of compounds of formula I.

Preferred compounds of formula IV, which are useful intermediates for the preparation of compounds of formula I, are:

(7S,8aS)-7-hydroxymethyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-
5 pyrrolo[1,2-a]pyrazine,

(7S,8aS)-7-hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-
octahydro-pyrrolo[1,2-a]pyrazine,

(7R,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-
pyrrolo[1,2-a]pyrazine,
10 (7R,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-
pyrrolo[1,2-a]pyrazine,

(7S,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-
pyrrolo[1,2-a]pyrazine, and

(7S,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-
15 pyrrolo[1,2-a]pyrazine.

This invention also relates to the pharmaceutically acceptable acid addition salts of compounds of the formula I. The compounds of formula I are basic in nature and are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid
20 addition salts of those compounds of formula I are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, fumarate, gluconate,
25 glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, and p-toluenesulfonate.

The term "one or more substituents", as used herein, includes from one to the maximum number of substituents possible based on the number of available bonding sites.

30 The chemist of ordinary skill will recognize that certain combinations of substituents may be chemically unstable and will avoid these combinations or alternatively protect sensitive groups with well known protecting groups.

-12-

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

5 The term "alkoxy", as used herein, unless otherwise indicated, refers to radicals having the formula -O-alkyl, wherein "alkyl" is defined as above.

The compounds of formula I contain chiral centers and therefore exist in different enantiomeric forms. This invention relates to all stereoisomers of compounds of the formula I and mixtures thereof.

10 The term "disorders of the dopamine system", as referred to herein, refers to disorders the treatment or prevention of which can be effected or facilitated by altering (i.e., increasing or decreasing) dopamine mediated neurotransmission.

The compounds in accordance with the present invention, being ligands for dopamine receptor subtypes, especially the dopamine D₄ receptor, within the body, are accordingly of use in the treatment or prevention of disorders of the dopamine
15 system.

It is generally accepted knowledge that dopamine receptors are important for many functions in the animal body. For example, altered functions of these receptors participate in the genesis of psychosis, addiction, sleep, feeding, learning, memory, sexual behavior, and blood pressure.

20 This invention provides dopamine ligands that are of use in the treatment and/or prevention of disorders of the dopamine system, including schizophrenia, nausea, Parkinson's disease, tardive dyskinesias and extrapyramidal side-effects associated with treatment by conventional neuroleptic agents, neuroleptic malignant syndrome, and disorders of hypothalamic-pituitary function such as
25 hyperprolactinaemia and amenorrhoea.

Upper gastrointestinal tract motility is believed to be under the control of the dopamine system. The compounds according to the present invention are of use in the prevention or treatment of gastrointestinal disorders, and the facilitation of gastric emptying.

30 Dependence-inducing agents such as cocaine and amphetamine have been shown to interact with the dopamine system. Compounds capable of counteracting this effect, including the compounds in accordance with the present invention, are

-13-

accordingly of value in the prevention or reduction of dependence on a dependence-inducing agent.

Dopamine is known to be a peripheral vasodilator; for example, it has been shown to exert a dilatory effect on the renal vascular bed. This shows that the compounds of the present invention are beneficial in controlling vascular blood flow.

The localization of dopamine receptor mRNA in rat heart and large vessels has been noted. This suggests a role for dopamine receptor ligands in controlling cardiovascular function, either by affecting cardiac and smooth muscle contractility or by modulating the secretion of vasoactive substances. The compounds according to the present invention are of assistance in the prevention or treatment of such conditions as hypertension and congestive heart failure.

The presence of D4 receptor mRNA in rat retina has been noted (Cohen, et al. *Proc. Nat. Acad. Sci.*, 1992, 89, 12093), suggesting that dopamine and D4 receptors play a role in ocular function. The compounds of this invention may therefore be useful in the treatment of ocular disorders. Furthermore, D4 receptors influence melatonin biosynthesis in chick retina (Zawilska, Nowak, *Neuroscience Lett.*, 1994, 166, 203), and since melatonin has been used for the treatment of sleep disorders, the compounds of this invention may be useful for the treatment of sleep disorders as well.

Detailed Description of the Invention

The compounds of formula I are readily prepared by methods which are summarized in the Schemes.

While the overall routes and various intermediates in the Schemes are novel, the individual chemical steps are generally analogous to known chemical transformations. Generally suitable conditions are found in the prior art. Isolation and purification of the products is accomplished by standard procedures which are known to a chemist of ordinary skill. Particularly well-suited conditions are exemplified below.

As used herein, the expression "reaction inert solvent" refers to a solvent system in which the components do not interact with starting materials, reagents, intermediates of products in a manner which adversely affects the yield of the desired product.

During any of the following synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned.

-14-

This may be achieved by means of conventional protecting groups, such as those described in T. W. Greene, *Protective Groups in Organic Chemistry*, John Wiley & Sons, 1981; and T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Chemistry*, John Wiley & Sons, 1991.

5 The expression "nitrogen protecting group" as used herein means a moiety which when coupled with a basic nitrogen will remain inert while other reactions are carried out. The nitrogen protecting group may then be removed under mild conditions yielding the free amino group. This invention contemplates two types of nitrogen protecting groups: those which may be removed by treatment with strong
10 acid and those which may be removed by hydrogenation.

 Examples of nitrogen protecting groups removed by strong acid are tert-butoxycarbonyl, methoxycarbonyl, ethoxycarbonyl, trimethylsilyloxycarbonyl, 1-adamantyloxycarbonyl, vinyloxycarbonyl, diphenylmethoxycarbonyl, trityl, acetyl and benzoyl.

15 Examples of nitrogen protecting groups removed by hydrogenation are benzyloxycarbonyl, 9-fluorenylmethyloxycarbonyl, 2-phenylethyloxycarbonyl, benzyl, p-methoxybenzyloxycarbonyl, and p-nitrobenzyloxycarbonyl. A preferred nitrogen protecting group is benzyl.

 Compounds of formula VII are useful intermediates for the preparation of
20 compounds of formula I. Compounds of formula VII wherein n is 0 are known (Diafi, L.; et al. *J. Het. Chem.*, 1990, 27, 2181). Compounds of formula VII wherein n is 1 may be prepared according to Scheme 1. Compounds of formula VII wherein n is 2 may be prepared according to Scheme 2.

 Scheme 1 summarizes methods suitable for preparing compounds VII wherein
25 n is 1. Compounds of the formulas V and VI wherein R₂ is hydrogen or methyl are known (Jones, R. C. F.; Howard, K. J. *J. Chem. Soc. Perkin Trans. 1*, 1993, 2391), and additional examples of VII wherein R₂ is a straight chain, branched or cyclic C₁ to C₆ alkyl may be prepared by this method using acrylate esters where the R₂ substituent is placed at the 2- position of the acrylate starting material. Compounds VI
30 may be converted into VII by the action of a strong hydride reducing agent in a reaction inert solvent, preferably an ether such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 65 °C. Many suitable reducing agents are known,

-15-

particularly hydrides of aluminum and boron, and lithium aluminum hydride or diborane are preferred reducing agents. Compounds VII prepared by this method may be produced as a mixture of stereoisomers, and the stereoisomers are separated by chromatographic methods known to a chemist of ordinary skill.

5 Compound V wherein R₂ is hydrogen may be used to prepare VIII by the action of a base in a lower alcohol solvent such as methanol, ethanol or propanol with an amine base, wherein 1,8-diazbicyclo[5.4.0]undec-7-ene (DBU) or 1,5-diazbicyclo[4.3.0]non-5-ene (DBN) are preferred bases, at a temperature from about 0 °C to about 100 °C, where a preferred temperature is from about 20 °C to about 60
10 °C. Reduction of VIII to IX may be accomplished by exposing a mixture of VIII and a noble metal catalyst, wherein palladium is a preferred noble metal catalyst, wherein the metal may be conveniently suspended on an inert solid support such as charcoal, in a solvent such as ethyl acetate or hexane or a mixture thereof, to an atmosphere of hydrogen gas at a pressure of about 1 to 100 atmospheres, where a preferred
15 pressure of hydrogen gas is about one to about ten atmospheres. Compound IX may be converted into VII by the action of a strong hydride reducing agent in a reaction inert solvent, preferably an ether such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 65 °C. Many suitable reducing agents are known, particularly
20 hydrides of aluminum and boron, and lithium aluminum hydride or diborane are preferred reducing agents. Compounds VII prepared by this method may be produced as a mixture of stereoisomers, and the stereoisomers are separated by chromatographic methods known to a chemist of ordinary skill.

 Compound IX may be converted into compound VI by treating a solution of IX
25 in an ether solvent, preferably tetrahydrofuran (THF), with an alkali metal amide base, preferably lithium diisopropylamide or potassium hexamethyl disilazide, at a temperature from about 0 °C to about -100 °C, preferably from about -50 °C to about -80 °C, followed by addition of an appropriate alkylating agent, such as an alkyl halide, wherein alkyl halides of one to six carbon atoms and bromide or iodide are
30 preferred. Compound IX prepared by this method may be converted into VII by the action of a strong hydride reducing agent in a reaction inert solvent, preferably an ether such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about

-16-

65 °C. Many suitable reducing agents are known, particularly hydrides of aluminum and boron, and lithium aluminum hydride or diborane are preferred reducing agents.

Referring to Scheme 2, compounds of formula VII wherein n is 2 may be prepared from compounds VII where n is 1 by first treating compound VII (n=1) with
5 an alkyl- or aryl-sulfonyl chloride, where methane sulfonyl chloride or tolyl sulfonyl chloride are preferred, in a reaction inert solvent, where methylene chloride or chloroform are preferred, at a temperature from about -10 °C to about 20 °C in the presence of a trialkyl amine base, where triethyl amine or diisopropyl ethyl amine are preferred. Treatment of the alkyl- or aryl-sulfonyl ester thus formed with an alkali
10 metal salt of cyanide, where potassium cyanide is preferred, in a polar solvent, where dimethylformamide (DMF) or dimethyl sulfoxide (DMSO) are preferred, at a temperature from about 20 °C to about 100 °C, where 50 °C to 80 °C is preferred, is a method that may be used to produce X. Conversion of nitrile X into aldehyde XI may be accomplished by treating a solution of X in an ether solvent, preferably
15 tetrahydrofuran (THF), with a hydride reducing agent, preferably diisobutyl aluminum hydride, at a temperature from about 0 °C to about 80 °C, preferably from about 20 °C to about 60 °C. Many reagents capable of reducing aldehyde XI to alcohol VII where n is 2 are known, and this transformation may be conveniently accomplished by using a hydride of boron, preferably sodium borohydride or sodium
20 cyanoborohydride, in a lower alcohol solvent, preferably methanol or ethanol, at a temperature from about -10 °C to about 30 °C.

Scheme 3 summarizes methods for preparing compounds I from compounds VII wherein X is O, S, NR₄, C(=O)O, C(=O)NR₄ or CHR₄.

Referring to Scheme 3, a method for coupling alcohols VII with compounds of
25 the formula R₁-(CH₂)_m-XH wherein X is O, S, NR₄, C(=O)O or C(=O)NR₄ to produce XII involves conversion of the alcohol moiety of VII into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with alcohol VII in the presence of a trialkyl amine, preferably
30 triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to about 30 °C. In a second step, a compound of formula R₁-(CH₂)_m-XH wherein X is O, S, NR₄, C(=O)O or C(=O)NR₄, is combined with a suitable base such as an alkali metal hydride, preferably sodium hydride, or an

-17-

alkali metal carbonate, preferably sodium carbonate or potassium carbonate, and the alkyl- or arylsulfonyl derivative of VII in a polar reaction inert solvent, preferably dimethyl formamide (DMF) or dimethylsulfoxide (DMSO), at a temperature from about 0 °C to about 150 °C, preferably from about 50 °C to about 100 °C.

- 5 In the case wherein X is O or S and m is 0, compounds XII may be prepared from VII by combining approximately equimolar quantities of VII, an alcohol of formula R_1OH or a thiol of formula R_1SH , triarylphosphine, preferably triphenylphosphine (Ph_3P), and dialkyl azodicarboxylate, preferably diethyl azodicarboxylate (DEAD), in a relatively polar ether solvent such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from
10 about 25 °C to about 60 °C.

- Compounds XII wherein X is $C(=O)O$ may be conveniently prepared by reacting VII with a carboxylic acid halide of the formula $R_1(CH_2)_mC(=O)Y$, wherein Y is a halogen atom, preferably chloro, in a reaction inert solvent, preferably methylene
15 chloride or chloroform, accompanied by a mild base, preferably triethylamine, 4-(dimethylamino)pyridine, or pyridine, at a temperature from about -10 °C to about 35 °C. In another preferred method, compound XII may be prepared by combining VII with a compound of the formula $R_1(CH_2)_mCO_2H$, a triarylphosphine, preferably triphenylphosphine (Ph_3P), and dialkyl azodicarboxylate, preferably diethyl
20 azodicarboxylate (DEAD), in a relatively polar ether solvent, preferably tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane, at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C.


- In the case wherein X is O and m is 1 or 2, XII may be conveniently prepared by reacting VII with a compound of formula $R_1-(CH_2)_m-L$ wherein L is a leaving group
25 and a suitable base such as an alkali metal hydride, preferably sodium hydride, or an alkali metal carbonate, preferably sodium carbonate or potassium carbonate, in a reaction inert solvent, preferably a polar solvent such as dimethylformamide (DMF), dimethyl sulfoxide (DMSO) or acetonitrile, at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C. The term "leaving group" refers
30 to groups which may be replaced by other groups under suitable conditions, and includes, for example, halogen, lower-alkylsulfonyl and arylsulfonyl. Methanesulfonyl, bromo or iodo are preferred leaving groups.

-18-

A method of preparing compounds of formula XII wherein X is CHR_4 from VII involves conversion of the alcohol moiety of VII into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with VII in the presence of a trialkyl amine, preferably triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10°C to about 30°C . In a second step, the alkyl- or arylsulfonyl derivative is combined with a compound of the formula $\text{R}_1(\text{CH}_2)_m\text{CH}(\text{R}_4)\text{MgY}$ wherein Y is a halogen atom, preferably chloro, bromo, or iodo, and a salt of copper(I), preferably cuprous bromide or cuprous iodide, in an ether solvent, preferably tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane, at a temperature from about -100°C to about 50°C , preferably from about -80°C to about 25°C .

Compounds of formula XII may be converted into compounds XIII by combining XII with a noble metal catalyst on an inert solid material, preferably palladium on charcoal, in a polar solvent such as a lower alcohol, preferably methanol or ethanol, to an atmosphere of hydrogen gas at a pressure of about one to twenty atmospheres and a temperature from about 0°C to about 100°C , preferably from about 25°C to about 60°C . Alternately, XIII may be conveniently prepared from XII by the action of ammonium formate and a noble metal catalyst, preferably palladium on charcoal, in a polar solvent such as water or a lower alcohol, preferably methanol or ethanol, or a mixture of water and a lower alcohol, at a temperature from about 0°C to about 100°C , preferably from about 25°C to about 60°C .

Compounds of formula I may be prepared from compounds of formula XIII using methods which are directly analogous to those described by Bright and Desai (US Pat. 5,122,525). The term "activated form of R_3 " means a chemical derivative of R_3 having the formula



-19-

- wherein L is a leaving group. The term "leaving group" (L) refers to groups which may be replaced by other groups under suitable conditions, and include, for example, halogen, lower-alkylsulfonyl and arylsulfonyl. Activated forms of R₃ may also be
- 5 derivatives of benzene bearing an electron withdrawing group (EWG) and a leaving group (L) in the ortho- or para- positions relative to one other:



- 10 Where the activated form of R₃ is a derivative of benzene, halogens are the preferred leaving groups, especially fluoro, and nitro or cyano are examples of preferred electron withdrawing groups. The reaction of compound XIII with an activated form of R₃ to form I is conveniently carried out in reaction inert solvents such as water, C₁-C₆ alcohols, or dimethyl sulfoxide, wherein water, propanol,
- 15 butanol, or pentanol are preferred, at a temperature from about 30 °C to about 170 °C, preferably from about 60 °C to about 110 °C. The presence of an acid acceptor such as a trialkyl amine or an alkali carbonate may be useful, and sodium carbonate or potassium carbonate is preferred.

- Compounds of formula I wherein X is O, S, NR₄, C(=O)O, C(=O)NR₄ or CHR₄
- 20 may also be prepared from VII by first removing the benzyl nitrogen protecting group. The formation of amino alcohols XIV from VII by this method may be conveniently carried out by combining VII with a noble metal catalyst on an inert solid material, preferably palladium on charcoal, in a polar solvent such as a lower alcohol, preferably methanol or ethanol, to an atmosphere of hydrogen gas at a pressure of
- 25 about one to twenty atmospheres and a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C. Alternately, XIV may be conveniently prepared from VII by the action of ammonium formate and a noble metal catalyst, preferably palladium on charcoal, in a polar solvent such as water or a lower alcohol, preferably methanol or ethanol, or a mixture of water and a lower alcohol, at a

-20-

temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C.

Compounds XV may be prepared by reacting XIV with an activated form of R₃ in a reaction inert solvent such as water, C₁-C₆ alcohols, or dimethyl sulfoxide, wherein water, propanol, butanol, or pentanol are preferred, at a temperature from about 30 °C to about 170 °C, preferably from about 60 °C to about 110 °C. The presence of an acid acceptor such as a trialkyl amine or an alkali carbonate may be useful, and sodium carbonate or potassium carbonate are preferred.

A useful method for coupling alcohols XV with compounds of the formula R₁-(CH₂)_m-XH wherein X is O, S, NR₄, (C=O)O or C(=O)NR₄ to produce I involves conversion of the alcohol moiety of XV into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with alcohol XV in the presence of a trialkyl amine, preferably triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to about 30 °C. In a second step, a compound of formula R₁-(CH₂)_m-XH wherein X is O, S, NR₄, (C=O)O or C(=O)NR₄ is combined with a suitable base such as an alkali metal hydride, preferably sodium hydride, or an alkali metal carbonate, preferably sodium carbonate or potassium carbonate, and the alkyl- or arylsulfonyl derivative of XV in a polar reaction inert solvent, preferably dimethyl formamide (DMF) or dimethylsulfoxide (DMSO), at a temperature from about 0 °C to about 150 °C, preferably from about 50 °C to about 100 °C.

In the case wherein X is O or S and m is 0, compounds I may be prepared from XV by combining approximately equimolar quantities of XV, an alcohol of formula R₁OH or a thiol of formula R₁SH, triarylphosphine, preferably triphenylphosphine (Ph₃P), and dialkyl azodicarboxylate, preferably diethyl azodicarboxylate (DEAD), in a relatively polar ether solvent such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C.

Compounds I wherein X is O and m is 1 or 2, may be conveniently prepared by reacting XV with a compound of formula R₁-(CH₂)_m-L wherein L is a leaving group and a suitable base such as an alkali metal hydride, preferably sodium hydride,

-21-

or an alkali metal carbonate, preferably sodium carbonate or potassium carbonate, in a reaction inert solvent, preferably a polar solvent such as dimethylformamide (DMF), dimethyl sulfoxide (DMSO) or acetonitrile at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C. The term "leaving group" refers to groups which may be replaced by other groups under suitable conditions, and include, for example, halogen, lower-alkylsulfonyl and arylsulfonyl. Methanesulfonyl, bromo or iodo are preferred leaving groups.

Compounds I wherein X is C(=O)O may be conveniently prepared by reacting XV with a carboxylic acid halide of the formula $R_1(CH_2)_mC(=O)Y$, wherein Y is a halogen atom, preferably chloro, in a reaction inert solvent, preferably methylene chloride or chloroform, accompanied by a mild base, preferably triethylamine, 4-(dimethylamino)pyridine, or pyridine, at a temperature from about -10 °C to about 35 °C. In another preferred method, compound I may be prepared by combining XV with a compound of the formula $R_1(CH_2)_mCO_2H$, a triarylphosphine, preferably triphenylphosphine (Ph_3P), and dialkyl azodicarboxylate, preferably diethyl azodicarboxylate (DEAD), in a relatively polar ether solvent, preferably tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane, at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C.

Compounds I wherein X is C(=O)N(R₄) may be conveniently prepared by conversion of the alcohol moiety of XV into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with XV in the presence of a trialkyl amine, preferably triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to about 30 °C. In a second step, the alkyl- or arylsulfonyl derivative is combined with a compound of the formula $R_1(CH_2)_mC(=O)NHR_4$, and a base, preferably sodium hydride or potassium carbonate, in a polar solvent, preferably dimethyl formamide (DMF) or dimethyl sulfoxide (DMSO), at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C. In cases wherein R₄ is hydrogen, treating compound I wherein X is C(=O)NH with a compound of formula R₄-L, wherein L is a leaving group, preferably chloro, bromo, iodo, lower-alkylsulfonyl ester or arylsulfonyl ester, and a base, preferably sodium

-22-

hydride, in a polar reaction inert solvent, preferably dimethyl formamide (DMF) or dimethyl sulfoxide (DMSO), at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C, is another preferred method for preparing compounds of formula I wherein X is C(=O)N(R₄).

- 5 A method of preparing compounds of formula I wherein X is CHR₄ from XV involves conversion of the alcohol moiety of XV into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with XV in the presence of a trialkyl amine, preferably triethylamine, in a
- 10 reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to about 30 °C. In a second step, the alkyl- or arylsulfonyl derivative is combined with a compound of the formula R₁(CH₂)_mCH(R₄)MgY wherein Y is a halogen atom, preferably chloro, bromo, or iodo, and a salt of copper(I), preferably cuprous bromide or cuprous iodide, in an ether solvent,
- 15 preferably tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane, at a temperature from about -100 °C to about 50 °C, preferably from about -80 °C to about 25 °C.

Scheme 4 summarizes methods suitable for preparing compounds I wherein X is C(=O), CH(OH) or CHR₄.

- 20 A method for preparing compounds XVI from XV involves conversion of the alcohol moiety of XV into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with XV in the presence of a trialkyl amine, preferably triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to
- 25 about 30 °C. In a second step, the alkyl- or arylsulfonyl derivative is combined with an alkali metal cyanide, preferably sodium cyanide or potassium cyanide, in a polar, reaction inert solvent, preferably dimethylformamide (DMF) or dimethyl sulfoxide (DMSO), at a temperature from about 25 °C to about 100 °C, preferably from about 35 °C to about 70 °C.

- 30 Compounds of formula I wherein X is C(=O) may be prepared by treating XVI with a compound of the formula R₁(CH₂)_m-MgY wherein Y is a halogen atom, preferably chloro, bromo or iodo, in an ether solvent, preferably tetrahydrofuran (THF) or diethyl ether, at a temperature from about -30 °C to about 60 °C, preferably from

-23-

about -10 °C to about 30 °C. Hydrolysis of the reaction mixture in a second step with a dilute aqueous solution of a strong inorganic acid, preferably hydrochloric acid or sulfuric acid, may be beneficial.

Many reagents capable of reducing compounds I wherein X is C(=O) to compounds I wherein X is CH(OH) are known, and this transformation may be conveniently accomplished by using a hydride of boron, preferably sodium borohydride, in a lower alcohol solvent, preferably methanol or ethanol, at a temperature from about -10 °C to about 30 °C.

A method for preparing compounds I wherein X is CH(R₄) from I wherein X is CH(OH) involves conversion of the alcohol moiety of I wherein X is CH(OH) into a leaving group such as lower-alkylsulfonyl ester or arylsulfonyl ester as the first step of a two-step process. The alkyl- or arylsulfonyl ester is first prepared by the reaction of an alkyl- or arylsulfonyl chloride with I wherein X is CH(OH) in the presence of a trialkyl amine, preferably triethylamine, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10 °C to about 30 °C. In a second step, the alkyl- or arylsulfonyl derivative is combined with a compound of the formula R₄-MgY wherein Y is a halogen atom, preferably chloro, bromo, or iodo, and a salt of copper(I), preferably cuprous bromide or cuprous iodide, in an ether solvent, preferably tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane, at a temperature from about -100 °C to about 50 °C, preferably from about -80 °C to about 25 °C.

Scheme 5 summarizes methods suitable for preparing compounds of formula I wherein X is OC(=O) or N(R₄)C(=O).

Compounds XVII may be prepared by treating XVI with an aqueous solution of a strong inorganic acid, preferably hydrochloric acid or sulfuric acid, at a temperature from about 20 °C to about 120 °C, preferably from about 50 °C to about 100 °C.

Many methods are known which are suitable for coupling carboxylic acids of formula XVII with alcohols of the formula R₁(CH₂)_m-OH to form compounds I wherein X is OC(=O). In a preferred method, a carboxylic acid of formula XVII, an alcohol of formula R₁(CH₂)_m-OH, a tertiary amine, preferably triethyl amine, diisopropyl ethyl amine or 4-(dimethylamino)pyridine, and 1-hydroxybenzotriazole hydrate (HOBt) dissolved in a reaction inert solvent, preferably methylene chloride or chloroform, is treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DEC) at a

-24-

temperature from about 0 °C to about 50 °C, preferably from about 10 °C to about 30 °C.

Many methods are known which are suitable for coupling carboxylic acids of formula XVII with amines of the formula $R_1(CH_2)_m-NHR_4$ to form compounds I wherein X is $N(R_4)C(=O)$. In a preferred method, a carboxylic acid of formula XVII, an amine of formula $R_1(CH_2)_m-NHR_4$, a tertiary amine, preferably triethyl amine or diisopropyl ethyl amine, and 1-hydroxybenzotriazole hydrate (HOBT) dissolved in a reaction inert solvent, preferably methylene chloride or chloroform, is treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DEC) at a temperature from about 0 °C to about 50 °C, preferably from about 10 °C to about 30 °C. In cases wherein R_4 is hydrogen, treating compound I wherein X is $NHC(=O)$ with a compound of formula R_4-L , wherein L is a leaving group, preferably chloro, bromo, iodo, lower-alkylsulfonyl ester or arylsulfonyl ester, and a base, preferably sodium hydride, in a polar reaction inert solvent, preferably dimethyl formamide (DMF) or dimethyl sulfoxide (DMSO), at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C, is another preferred method for preparing compounds of formula I wherein X is $N(R_4)C(=O)$.

Scheme 6 summarized methods for preparing compound XIV wherein n is 1 and R_2 is hydrogen.

Methods for preparing the stereoisomers of compound XVIII have been reported by Bridges, et al. (*J. Med. Chem.*, 1991, 34, 717). The formation of amine XIX from XVIII may be conveniently carried out by combining XVIII with a noble metal catalyst on an inert solid material, preferably palladium on charcoal, in a polar solvent such as a lower alcohol, preferably methanol or ethanol, under an atmosphere of hydrogen gas at a pressure of about one to twenty atmospheres and a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C. Alternately, XIX may be conveniently prepared from XVIII by the action of ammonium formate and a noble metal catalyst, preferably palladium on charcoal, in a polar solvent such as water or a lower alcohol, preferably methanol or ethanol, or a mixture of water and a lower alcohol, at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 60 °C.

-25-

A preferred method for preparing compound XX from compound XIX involves combining 2-(phthalimido)acetaldehyde with compound XIX, an alkali metal salt of a carboxylic acid, preferably sodium acetate or sodium propionate, and a boron hydride reducing agent, preferably sodium (triacetoxy)borohydride, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about 5 °C to about 30 °C in the first step of a two step process. In the second step, the 2-(phthalimido)ethyl derivative thus formed is combined with a primary amine capable of causing the removal of the phthalimido protecting group, preferably hydrazine hydrate or methyl amine, in water or a lower alcohol solvent, preferably water, methanol, or ethanol, or a mixture thereof, at a temperature from about 0 °C to about 60 °C, preferably from about 5 °C to about 30 °C.


Compound XX is converted into XIV wherein n is 1 by the action of a strong hydride reducing agent in a reaction inert solvent, preferably an ether such as tetrahydrofuran (THF), 1,4-dioxane or 1,2-dimethoxyethane at a temperature from about 0 °C to about 100 °C, preferably from about 25 °C to about 65 °C. Many suitable reducing agents are known, particularly hydrides of aluminum and boron, and lithium aluminum hydride or diborane are preferred reducing agents.

Compounds I wherein X is S may be converted into compounds I wherein X is SO by the action with an oxidizing agent, preferably about one molar equivalent of m-chloroperoxybenzoic acid, in a reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -100 °C to about 0 °C, preferably from about -80 °C to about -20 °C.

Compounds I where X is S or SO may be converted into compounds I where X is SO₂ by the action with an oxidizing agent, preferably m-chloroperoxybenzoic acid, in reaction inert solvent, preferably methylene chloride or chloroform, at a temperature from about -10°C to about 50 °C, preferably from about 0 °C to about 30 °C.


-26-

Scheme 1.




5

Scheme 2.




-27-

Scheme 3.




-28-

Scheme 4.




5.

Scheme 5.



-29-

Scheme 6.



5 The novel compounds of the formula I and the pharmaceutically acceptable salts thereof (herein "the therapeutic compounds of this invention") are useful as dopaminergic agents, i.e., they possess the ability to alter dopamine mediated neurotransmission in mammals, including humans. They are therefore able to function as therapeutic agents in the treatment of a variety of conditions in mammals, the treatment or prevention of which can be effected or facilitated by an increase or decrease in dopamine mediated neurotransmission.

10 The compounds of the formula I that are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

25 The therapeutic compounds of this invention can be administered orally, transdermally (e.g. through the use of a patch), parenterally or topically. Oral

-30-

administration is preferred. In general, these compounds are most desirably administered in dosages ranging from about 0.1 mg up to about 1000 mg per day, or 1 mg to 1000 mg per day in some cases, although variations may occur depending on the weight and condition of the person being treated and the particular route of administration chosen. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

The therapeutic compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the two routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the novel therapeutic compounds of this invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, for example. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

-31-

For parenteral administration, solutions of a compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

Dopaminergic activity is related to the ability of compounds to bind to mammalian dopamine receptors, and the relative ability of compounds of this invention to inhibit [³H]-spiperone binding to human dopamine D₄ receptor subtypes expressed in clonal cell lines was measured using the following procedure.

The determination of D₄ receptor binding ability has been described by Van Tol, et al. (*Nature*, 1991, 350, 610). Clonal cell lines expressing the human dopamine D₄ receptor are harvested and homogenized (polytron) in a 50 mM Tris:HCl (pH 7.4 at 4 °C) buffer containing 5 mM EDTA, 1.5 mM calcium chloride (CaCl₂), 5 mM magnesium chloride (MgCl₂), 5 mM potassium chloride (KCl) and 120 mM sodium chloride (NaCl). The homogenates are centrifugated for 10-15 min. at 48,000 g, and the resulting pellets resuspended in a buffer at a concentration of 150-250 mg/ml. For saturation experiments, 0.75 ml aliquots of tissue homogenate are incubated in triplicate with increasing concentrations of [³H]-spiperone (70.3 Ci/mmol; 10-3000 pM final concentration) for 30-120 minutes at 22 °C in a total volume of 1 ml. For competition binding experiments, assays are initiated by the addition of 0.75 ml of membrane and incubated in duplicate with the indicated concentrations of competing ligands (10⁻¹⁴-10⁻³ M) and/or [³H]-spiperone (100-300 pM) for 60-120 min at 22°C. Assays are terminated by rapid filtration through a Brandell cell harvester and the filters subsequently monitored for tritium as described by Sunahara, R.K. et al. (*Nature*, 1990, 346, 76). For all experiments, specific [³H]spiperone binding is defined


-32-

as that inhibited by 1-10 mM (+)-butaclamol. Binding data are analyzed by non-linear least square curve-fitting. The compounds of the Examples were tested in this assay, and all were found to have binding affinities (K_i) for the displacement of [^3H]-spiperone of less than 2 micromolar.

5 The following Examples are provided solely for the purposes of illustration and do not limit the invention which is defined by the claims.

Example 1

(7RS,8aSR)-7-(4-Fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



10

15

A solution of 2.00 g (5.9 mmol) of (7RS,8aSR)-7-(4-fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 2) and 4.1 mL (20.6 mmol) of 5M aqueous ammonium formate in 50 mL methanol was treated with an aqueous slurry of 0.200 g of 10% Pd/C. The reaction was refluxed for

20 48 hrs. The mixture was filtered and the solvent removed *in vacuo* to give an oily residue. The crude (7RS,8aSR)-7-(4-fluorophenoxy)methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine was combined with 1.2 g (9.1 mmol) of 2-chloro-5-fluoropyrimidine (Dunaiskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), 2.0 g (9.1 mmol) of sodium carbonate in 100 mL of water, and the mixture was gently

25 refluxed for 16 hrs. After cooling to room temperature, the mixture was extracted with methylene chloride (3x). The combined organic layers were dried (magnesium sulfate), filtered, and evaporated. Purification by flash silica gel chromatography eluting with 90:10 ethyl acetate:hexane gave 0.744 g (27%) of the title compound. mp


30 (-HCl) 120-122 °C. ^{13}C NMR (base, CDCl_3): δ 31.4, 35.3, 43.8, 49.0, 51.5, 57.6, 61.5, 71.4, 115.39, 115.50, 115.62, 115.92, 144.96, 145.24, 149.9, 153.2, 155.1, 155.7, 158.8. Anal calcd for $\text{C}_{18}\text{H}_{20}\text{F}_2\text{N}_4\text{O}$: C, 62.41; H, 5.82; N, 16.18. Found: C, 62.05, H, 5.99; N, 16.33.

-33-

Example 2

(7SR,8aSR)-7-(4-Fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




- 10 A solution of 0.870 g (2.6 mmol) of (7SR,8aSR)-7-(4-fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 2) and 1.8 mL (8.9 mmol) of 5M aqueous ammonium formate in 50 mL methanol was treated with an aqueous slurry of 0.100 g of 10% Pd/C. The reaction was refluxed for 24 h. The mixture was filtered and the solvent removed *in vacuo* to give an oily residue.
- 15 The crude (7SR,8aSR)-7-(4-fluorophenoxy)methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine was combined with 0.373 g (2.8 mmol) of 2-chloro-5-fluoropyrimidine (Dunaïskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), 0.650 g (6.1 mmol) of sodium carbonate in 50 mL of water, and the mixture was gently refluxed for 16 hrs. After cooling to room temperature, the mixture was extracted with
- 20 methylene chloride (3x). The combined organic layers were dried (magnesium sulfate), filtered, and evaporated. Purification by silica gel flash chromatography eluting with 85:15 ethyl acetate:hexane gave 0.444 g (50%) of the title compound. mp (·HCl) 211-213 °C. ¹³C NMR (base, CDCl₃): δ 31.7, 35.2, 43.8, 49.1, 51.4, 56.6, 62.3, 72.5, 115.45, 115.56, 115.89, 144.95, 145.24, 149.9, 153.2, 155.1, 155.6, 158.79,
- 25 158.90. Anal calcd for C₁₈H₂₀F₂N₄O: C, 62.41; H, 5.82; N, 16.18. Found: C, 62.15, H, 5.99; N, 16.38.

-34-

Example 3

(7RS,8aSR)-7-(4-Fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-7-methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5



10 A solution of 0.745 g (2.80 mmol) of (7RS,8aSR)-7-hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-7-methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 5) and 0.43 mL (3.08 mmol) of triethylamine in 30 mL of dry methylene chloride was chilled to 0 °C, and treated with methanesulfonyl chloride (0.228 mL, 2.94 mmol) in 15 mL of dry methylene chloride. After 1 h, the solution was washed
15 with water (2x), dried (magnesium sulfate), filtered and evaporated to give 0.915g (95%) of mesylate as a pale yellow solid.


A solution of 0.23 g (2.0 mmol) of 4-fluorophenol in 10 mL of dry DMF was treated with 0.096 g (2.4 mmol) of sodium hydride (60% oil dispersion), and the mixture heated at 50 °C for 1 h. A solution of 0.25 g (0.73 mmol) of mesylate in 10 mL
20 of dry DMF was added and the solution heated at 100 °C for 72 h. The mixture was cooled to room temperature, diluted with water, and extracted with diethyl ether (2x). The organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography eluting with 2:1 petroleum ether:diethyl ether gave 0.15 g (58%) of the title compound. mp (·HCl) 158-160 °C. ¹³C NMR
25 (base, CDCl₃): δ 26.4, 39.9, 40.6, 43.8, 49.1, 51.6, 62.8, 64.5, 76.0, 115.3, 115.5, 115.6, 115.9, 145.0, 145.2, 149.9, 153.2, 155.4, 155.7, 158.8, 158.9. HRMS calcd for C₁₉H₂₃F₂N₄O (MH⁺): 361.1840; found: 361.1861.

-35-

Example 4

(7RS,8aSR)-7-Phenoxymethyl-2-(5-fluoropyrimidin-2-yl)-7-methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5



10 A solution of 0.745 g (2.80 mmol) of (7RS,8aSR)-7-hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-7-methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 5) and 0.43 mL (3.08 mmol) of triethylamine in 30 mL of dry methylene chloride was chilled to 0 °C, and treated with methanesulfonyl chloride (0.228 mL, 2.94 mmol) in 15 mL of dry methylene chloride. After 1 h, the solution was washed
15 with water (2x), dried (magnesium sulfate), filtered and evaporated to give 0.915 g (95%) of mesylate as a pale yellow solid.


A solution of 0.19 g (2.0 mmol) of phenol in 10 mL of dry DMF was treated with 0.096 g (2.4 mmol) of sodium hydride (60% oil dispersion), and the mixture heated at 50 °C for 1 h. A solution of 0.25 g (0.73 mmol) of mesylate in 10 mL of dry
20 DMF was added and the solution heated at 100 °C for 72 h. The mixture was cooled to room temperature, diluted with water, and extracted with diethyl ether (2x). The organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography eluting with petroleum ether:diethyl ether (2:1) gave 0.18 g (72%) of the title compound. mp (-HCl) 189-191 °C. ¹³C NMR (base, CDCl₃):
25 δ 26.4, 40.0, 40.6, 43.8, 49.1, 51.6, 62.8, 64.5, 75.3, 114.5, 120.7, 129.4, 145.0, 145.2, 149.9, 153.2, 158.9, 159.3. HRMS calcd for C₁₉H₂₃FN₄O (MH⁺): 343.1934; found: 343.1951.

-36-

Example 5

(7R,8aSR)-7-(4-Fluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




10 A mixture of 3.8 g (26 mmol) of 2,5-dichloropyridine, 1.3 g (12 mmol) of sodium carbonate, 1.3 g (5.2 mmol) of (7R,8aSR)-7-(4-fluorophenoxy)methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 2), and 20 mL of isoamyl alcohol was heated at reflux for 18 h. The solvent was evaporated, the residue taken up in water and ethyl acetate, and the pH adjusted to 11 with sodium
15 carbonate. The layers were separated and the organic phase was dried (magnesium sulfate), filtered, and evaporated. Purification by medium pressure silica gel chromatography with ethyl acetate gave 35 mg (2%) of the title compound. mp (\cdot HCl) 202-206 $^{\circ}$ C. HRMS calcd for $C_{19}H_{21}ClFN_3O$ (MH $^{+}$): 362.1435, found: 362.1451.

Example 6

20 (7S,8aS)-7-(4-Fluorophenoxy)-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

25



A solution of 0.971 g (4.18 mmol) of (7R,8aS)-7-hydroxy-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Diafi, L.; et al. *J. Het. Chem.*, 1990, 27, 2181), 0.703 g (6.27 mmol) of 4-fluorophenol and 1.32 g (5.02 mmol) of
30 triphenylphosphine in 20 mL of dry THF was treated with 0.79 mL (5.02 mmol) of diethyl azodicarboxylate and the solution stirred at room temperature for 21 h. Excess HCl (g) in diethyl ether was added, the precipitate was collected on a Büchner funnel, washing with ethyl acetate. The gummy residue was dissolved in a mixture of ethyl


-37-

acetate and aqueous ammonium hydroxide, the layers were separated, the aqueous phase was extracted with more ethyl acetate (2x), the combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by silica gel MPLC with 95:5 ethyl acetate:methanol gave 1.3 g (95%) of (7S,8aS)-7-(4-fluorophenoxy)-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

A solution of 0.83 g (2.5 mmol) of (7S,8aS)-7-(4-fluorophenoxy)-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine in 10 mL of methanol and 1.8 mL of aqueous ammonium formate (5 M) was treated with an aqueous slurry of 0.325 g of 10% palladium on carbon and the mixture was stirred at room temperature for 24 h. The mixture was filtered through Celite, evaporated, re-evaporated with another 100 mL of chloroform, dissolved in 100 mL of chloroform, dried (magnesium sulfate), filtered and evaporated. The crude amine (0.61 g, ca. 2.5 mmol), 0.50 g (3.75 mmol) of 2-chloro-5-fluoropyrimidine (Dunaïskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), 0.86 g (6.2 mmol) of potassium carbonate and 15 mL of 2-propanol were refluxed for 5.5 h. The mixture was cooled to room temperature, diluted with water, and extracted with ethyl acetate (3x). The combined organic phase was washed with water (2x) and brine (1x), dried (magnesium sulfate), filtered and evaporated. Purification by silica gel MPLC starting with 70:30 ethyl acetate:hexane and ramping to 50:50 ethyl acetate:hexane at 30 min, gave 0.30 g (36%) of the title compound. mp (-HCl) 90-95 °C. ¹³C NMR (base, CDCl₃): δ 36.6, 43.6, 48.7, 60.5, 62.1, 75.6, 115.67, 115.98, 116.08, 116.18, 144.97, 145.26, 149.9, 153.2, 153.8, 155.6, 158.8. HRMS calcd for C₁₇H₁₉F₂N₄O (MH⁺): 333.1527, found: 333.1556.

Example 7

(7R,8aS)-7-(4-Fluorobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine




-38-

A solution of 0.75 g (3.15 mmol) of (7R,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 6) and 0.79 mL (6.3 mmol) of 4-fluorobenzyl bromide in 30 mL of dry DMF was treated with 0.15 g (3.8 mmol) of sodium hydride (60% oil dispersion), and the mixture was heated at 100 °C for 18 h. The mixture was cool, diluted with water, and extracted with diethyl ether (5x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatograph with 50:50 ethyl acetate:hexane gave 0.090 g (8%) of the title compound. mp (\cdot HCl) 74-79 °C. ^{13}C NMR (base, CDCl_3): δ 35.6, 43.9, 48.9, 51.2, 60.4, 60.7, 70.7, 76.7, 115.1, 115.4, 129.35, 129.46, 133.9, 144.9, 145.2, 149.9, 153.2, 158.9, 164.0. HRMS calcd for $\text{C}_{18}\text{H}_{20}\text{F}_2\text{N}_4\text{O}$ (MH^+): 347.1683, found: 347.1671.

Example 8

(7S,8aS)-7-(4-Fluorobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine




A solution of 1.15 mL (10.2 mmol) of 4-fluorobenzyl alcohol in 35 mL of dry DMF was treated with 0.48 g (12 mmol) of sodium hydride (60% oil dispersion), and the mixture was stirred at 50 °C for 30 min. A solution of 1.15 g (3.64 mmol) of (7R,8aS)-7-methanesulfonyloxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 7) in 35 mL of dry DMF was added and the solution stirred at 100 °C for 18 h. The solution was cooled, diluted with water, and extracted with diethyl ether (2x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with ethyl acetate gave 0.25 g (20%) of the title compound. mp (\cdot D-(-)-tartrate) 76-81 °C. ^{13}C NMR (base, CDCl_3): δ 36.3, 43.6, 48.8, 51.3, 60.2, 61.9, 70.4, 76.6, 115.0, 115.3, 129.43, 129.53, 134.0, 144.9, 145.2, 149.9, 153.2, 158.9, 160.6, 163.9. HRMS calcd for $\text{C}_{18}\text{H}_{20}\text{F}_2\text{N}_4\text{O}$ (MH^+): 347.1683, found: 347.1706.

-39-

Example 9

(7S,8aS)-2-(5-Fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate

5



10

A solution of 2.0 g (8.4 mmol) of (7R,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 6), 1.54 g (12.6 mmol) of benzoic acid, 2.65 g (10.1 mmol) of triphenylphosphine, and 1.59 mL (10.1 mmol) of diethyl azodicarboxylate (DEAD) in 85 mL of THF was stirred at ambient temperature for 16 h. The solvent was evaporated, and flash silica gel

15

chromatography with 1:1 hexane:ethyl acetate gave 2.5 g of partially purified material. A second chromatography with 10:1 methylene chloride:methanol gave 1.68 g (59%) of the title compound. mp (\cdot HCl) 134-135.5 $^{\circ}$ C. 13 C NMR (base, CDCl_3): δ 36.1, 43.7, 48.7, 51.2, 60.4, 61.8, 73.2, 128.3, 129.7, 130.1, 133.0, 145.0, 145.3, 150.0, 153.1, 158.9, 166.7. HRMS calcd for $\text{C}_{18}\text{H}_{20}\text{FN}_4\text{O}$ (MH^+): 343.1570, found:


20

343.1585.

Example 10

(7S,8aS)-7-(4-Fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

25



30

A solution of 0.25 g (0.99 mmol) of (7S,8aS)-7-hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 13), 0.167 g (1.49 mmol) of 4-fluorophenol, 0.31 g (1.19 mmol) of triphenylphosphine and 0.19 mL (1.2 mmol) of diethyl azodicarboxylate (DEAD) in 10 mL of dry THF was


-40-

stirred at ambient temperature for 16 h. The solvent was evaporated, the residue was dissolved in chloroform and washed with 1M sodium hydroxide. The organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 1:1 hexane:ethyl acetate and a second chromatography with

5 3:1 hexane:ethyl acetate gave 0.185 g (54%) of the title compound. mp (\cdot HCl) 207.5-208 °C. ^{13}C NMR (base, CDCl_3): δ 31.7, 35.2, 43.8, 49.1, 51.4, 56.6, 62.3, 72.5, 115.5, 115.6, 115.9, 144.9, 145.2, 149.9, 153.2, 155.1, 155.6, 158.8, 158.9. m/z (MH $^{+}$) 347.

Examples 11

- 10 (7S,8aS)-7-(Substituted-phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazines



The following compounds were prepared from (7S,8aS)-7-hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 13) and the appropriate phenol according to the method described in Example 10.

- 20 11a. (7S,8aS)-7-(3-Cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 83-85 °C. HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{FN}_5\text{O}$ (MH $^{+}$): 354.1730, found: 354.1716.
- 11b. (7S,8aS)-7-(4-Cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 183-185 °C. HRMS calcd for $\text{C}_{19}\text{H}_{21}\text{FN}_5\text{O}$ (MH $^{+}$): 354.1730, found: 354.1719.
- 25 11c. (7S,8aS)-7-(3,5-Difluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 205-206 °C. HRMS calcd for $\text{C}_{18}\text{H}_{20}\text{F}_3\text{N}_4\text{O}$ (MH $^{+}$): 365.1589, found: 365.1592.
- 11d. (7S,8aS)-7-(2-Nitrophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 151-153 °C. HRMS calcd for $\text{C}_{18}\text{H}_{21}\text{FN}_5\text{O}_3$ (MH $^{+}$): 374.1628, found: 374.1638.
- 30

-41-


- 11e. (7S,8aS)-7-(3-Nitrophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 92-95 °C. HRMS calcd
for C₁₈H₂₁FN₅O₃ (MH⁺): 374.1628, found: 374.1647.
- 11f. (7S,8aS)-7-(4-Nitrophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
5 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 189-191 °C. HRMS calcd
for C₁₈H₂₁FN₅O₃ (MH⁺): 374.1628, found: 374.1647.
- 11g. (7S,8aS)-7-(3-(Trifluoromethyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: HRMS calcd for C₁₉H₂₁F₄N₄O
(MH⁺): 397.1651, found: 397.1642.
- 10 11h. (7S,8aS)-7-(4-(Cyanomethyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 65-67 °C. HRMS calcd
for C₂₀H₂₃FN₅O (MH⁺): 368.1887, found: 368.1898.
- 11i. (7S,8aS)-7-(4-((Methoxycarbonyl)methyl)phenoxy)methyl-2-(5-fluoropyri-
midin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 50-52 °C.
15 HRMS calcd for C₂₁H₂₅FN₄O₃ (MH⁺): 401.1989, found: 401.1965.
- 11j. (7S,8aS)-7-(3-(Methoxycarbonyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-
yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 74-79 °C. HRMS
calcd for C₂₀H₂₄FN₄O₃ (MH⁺): 387.1832, found: 387.1866.
- 11k. (7S,8aS)-7-(3-(Ethyne)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
20 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 73-76 °C. HRMS calcd
for C₂₀H₂₂FN₄O (MH⁺): 353.1778, found: 353.1805.
- 11l. (7S,8aS)-7-(3-(Ethoxy)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 169-170 °C. HRMS calcd
for C₂₀H₂₆FN₄O₂ (MH⁺): 373.2040, found: 373.2016.
- 25 11m. (7S,8aS)-7-(Phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-
octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 76-79 °C. HRMS calcd for C₁₈H₂₂FN₄O
(MH⁺): 329.1778, found: 329.1784.

-42-

Example 12

(7S,8aS)-7-(4-Fluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




10 A solution of 0.25 g (0.93 mmol) of (7S,8aS)-7-hydroxymethyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 14), 0.157 g (1.40 mmol) of 4-fluorophenol, 0.29 g (1.12 mmol) of triphenylphosphine and 0.18 mL (1.1 mmol) of diethyl azodicarboxylate (DEAD) in 10 mL of dry THF was stirred at ambient temperature for 16 h. The solvent was evaporated, the residue was dissolved in chloroform and washed with 1M sodium hydroxide. The organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 1:1 hexane:ethyl acetate gave 0.24 g (71%) of the title compound. mp (\cdot HCl) 221-224 °C. HRMS calcd for $C_{19}H_{22}ClFN_3O$ (MH $^{+}$): 362.1435, found: 362.1415.

20

Examples 13

(7S,8aS)-7-(Substituted-phenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazines

25



The following compounds were prepared from (7S,8aS)-7-hydroxymethyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 14) and the appropriate phenol according to the method described in Example 12.

30

-43-

13a. (7S,8aS)-7-(3-Cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 220-224 °C. HRMS calcd for $C_{20}H_{22}ClN_4O$ (MH⁺): 369.1482, found: 369.1472.


13b. (7S,8aS)-7-(4-Cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine. Purification by flash silica gel chromatography with ethyl ether. mp (\cdot HCl) 245 °C (dec). HRMS calcd for $C_{20}H_{22}ClN_4O$ (MH⁺): 369.1482, found: 369.1465.

13c. (7S,8aS)-7-(3,5-Difluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine. Purification by flash silica gel chromatography with 65:35 diethyl ether:petroleum ether. mp (\cdot HCl) 220 °C (dec). HRMS calcd for $C_{19}H_{21}ClF_2N_3O$ (MH⁺): 380.1341, found: 380.1309.

13d. (7S,8aS)-7-(4-(Methoxycarbonyl)methylphenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: mp (\cdot HCl) 186-189 °C. HRMS calcd for $C_{22}H_{27}ClN_3O_3$ (MH⁺): 416.1741, found: 416.1765.

Example 14

(7S,8aS)-7-(3-Cyanophenoxy)methyl-2-(5-cyanopyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



A solution of 1.0 g (6.4 mmol) of (7S,8aS)-7-hydroxymethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 13, Step A), 1.77 g (12.8 mmol) of 2-chloro-5-cyanopyridine and 2.71 g (25.6 mmol) of sodium carbonate in 50 mL of isoamyl alcohol was heated at reflux for 18 h. The mixture was cooled to room temperature, diluted with ethyl acetate and water, the pH was adjusted to 11 with sodium carbonate, the layers were separated, and the aqueous layer extracted with ethyl acetate. The combined organic phase was dried (magnesium sulfate), filtered and evaporated to give (7S,8aS)-7-(3-cyanophenoxy)methyl-2-(5-cyanopyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

-44-

The (7S,8aS)-7-hydroxymethyl-2-(5-cyanopyridin-2-yl)-1,2,3,4, 6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine, 1.6 g (13.4 mmol) of 3-cyanophenol, and 2.8 g (11 mmol) of triphenylphosphine were dissolved in 20 mL of dry THF, the solution was treated with 1.7 mL (11 mmol) of diethyl azodicarboxylate (DEAD), and the reaction


5 stirred at ambient temperature for 16 h. The solvent was evaporated, the residue taken up in ethyl acetate, and washed with 1M NaOH. The organic phase was extracted with 1M HCl (3x), and the aqueous acid was washed with ethyl acetate (1x). The aqueous phase was made basic with 1M NaOH, extracted with ethyl acetate (3x), and the combined organic phase dried (magnesium sulfate), filtered and

10 evaporated. Purification by flash silica gel chromatography with 75:25 ethyl acetate:hexane gave 0.603 g (19%) of the title compound. mp (\cdot HCl) 197-200 $^{\circ}$ C. HRMS calcd for $C_{21}H_{22}N_5O$ (MH $^{+}$): 360.1824, found: 360.1802.

Example 15

(7S,8aS)-7-(4-Fluorophenoxy)methyl-2-(5-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

15



A solution of 0.25 g (0.97 mmol) of (7S,8aS)-7-hydroxymethyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 16), 0.164 g (1.46 mmol) of 4-fluorophenol, 0.31 g (1.15 mmol) of triphenylphosphine in

25 10 mL of THF was treated with 0.18 mL (1.15 mmol) of diethyl azodicarboxylate and the solution stirred at ambient temperature for 16 h. The solvent was removed by rotary evaporation, and the residue partitioned between chloroform and 1M sodium hydroxide. The layers were separated, and the organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with


30 1:1 ethyl acetate:hexane gave 0.225 g (66%) of the title compound. mp (\cdot HCl) 81-85 $^{\circ}$ C. HRMS calcd for $C_{21}H_{23}FN_3O$ (MH $^{+}$): 352.1825, found: 352.1817.

-45-

Examples 16

(7S,8aS)-7-(Substituted-phenoxy)methyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5



The following compounds were prepared from (7S,8aS)-7-hydroxymethyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 16) and the appropriate phenol according to the method described in Example 15.

16a. (7S,8aS)-7-(3-Cyanophenoxy)methyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine. mp 114-118 °C. HRMS calcd for C₂₂H₂₃N₄O (MH⁺): 359.1872, found: 359.1877.


16b. (7S,8aS)-7-(4-Cyanophenoxy)methyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine. mp 128-135 °C. HRMS calcd for C₂₂H₂₃N₄O (MH⁺): 359.1872, found: 359.1879.

16c. (7S,8aS)-7-(3-Ethoxyphenoxy)methyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine. mp 94-98 °C. HRMS calcd for C₂₃H₂₈N₃O₂ (MH⁺): 378.2182, found: 378.2145.

Example 17

(7S,8aS)-2-(5-Chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate

25



30

A solution of 0.595 g (2.35 mmol) of (7R,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 17), 0.430 g (3.52 mmol) of benzoic acid and 0.738 g (2.81 mmol) of triphenylphosphine in 25 mL of dry THF was treated with 0.44 mL (2.8 mmol) of diethyl azodicarboxylate, and the mixture


-46-

stirred at ambient temperature for 16 h. The solvent was evaporated and the residue purified by flash silica gel chromatography with 10:1 methylene chloride:acetone. A second chromatography of the major fraction with the same system gave 0.63 g (75%) of the title compound. mp (\cdot HCl) 202-205 °C. HRMS calcd for $C_{19}H_{21}ClN_3O_2$ (MH⁺): 358.1322, found: 358.1320.

Example 18

(7S,8aS)-7-(4-Fluorobenzyl)oxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

10



15


A solution of 0.23 g (0.91 mmol) of (7S,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 18) in 10 mL of dry THF was treated with 40 mg (1.0 mmol) of sodium hydride (60% oil dispersion), followed by 0.125 mL (1.0 mmol) of 4-fluorobenzyl bromide and 17 mg (0.05 mmol) of tetra-n-butylammonium iodide. The mixture was stirred at ambient temperature for 16 h, and warmed to 50 °C for 4 h. The suspension was cooled, the solvent was evaporated, and the residue partitioned between ethyl acetate and water. The layers were separated, and the organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 95:5 chloroform: methanol gave 0.135 g (41%) of the title compound. mp (\cdot HCl) 165-168 °C. HRMS calcd for $C_{19}H_{22}ClFN_3O$ (MH⁺): 362.1435, found: 362.1451.

25

-47-

Example 19


(7S,8aS)-7-(3-Cyanobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



5
10 A solution of 0.60 g (2.5 mmol) of (7S,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 15) in 30 mL of THF was treated with 0.41 g (10 mmol) of sodium hydride (60% oil dispersion), followed by 0.75 g (3.8 mmol) of 3-cyanobenzyl bromide and 30 mg (0.1 mmol) of tetra-n-butylammonium iodide. The mixture was stirred 50 °C for 16 h, cooled to room
15 temperature, the solvent was evaporated, and the residue partitioned between ethyl acetate and water. The layers were separated, the organic phase was washed with water and brine, dried (magnesium sulfate), filtered and evaporated. Purification by MPLC silica gel chromatography with ethyl acetate gave 0.11 g (12%) of the title compound. mp (·HCl) 95-100 °C. HRMS calcd for C₁₉H₂₁FN₅O (MH⁺): 345.1730;
20 found: 345.1739.

Example 20

(7S,8aS)-7-(4-(2-Hydroxyethyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



25
30 A solution of 0.13 g (0.33 mmol) of (7S,8aS)-7-(4-((methoxycarbonyl)methyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Example 11i) in 15 mL of anhydrous ethyl ether


-48-

was added dropwise to an ice-cold suspension of 0.025 g (0.65 mmol) of lithium aluminum hydride in 15 mL of anhydrous ethyl ether and the mixture stirred for 30 min. The reaction was carefully quenched at 0 °C with 0.025 mL of water, 0.025 mL of 15% sodium hydroxide, and 0.075 mL of water. The precipitate which formed was
5 filtered through Celite, the filtrate concentrated *in vacuo*, and purification of the residue by flash silica gel chromatography with 95:5 ethyl acetate:methanol gave 0.075 g (63%) of the title compound. mp (\cdot HCl) 145-147 °C. HRMS calcd for $C_{20}H_{26}FN_4O_2$ (MH⁺): 373.2040, found: 373.2054.

Example 21

10 (7S,8aS)-7-(3-(Hydroxymethyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-
pyrrolo[1,2-a]pyrazine

15




A solution of 0.15 g (0.39 mmol) of (7S,8aS)-7-(3-(methoxycarbonyl)phenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Example 11j) in 15 mL of anhydrous ethyl ether was added dropwise to an ice-cold suspension of 0.029 g (0.78 mmol) of lithium aluminum hydride in 15 mL of anhydrous ethyl ether and the mixture stirred for 30 min. The reaction was carefully quenched at 0 °C with 0.029 mL of water, 0.029 mL of
25 15% sodium hydroxide, and 0.087 mL of water. The precipitate which formed was filtered through Celite, the filtrate concentrated *in vacuo*, and purification of the residue by flash silica gel chromatography with 95:5 chloroform:methanol gave 0.099 g (72%) of the title compound. mp (\cdot HCl) 85-89 °C. HRMS calcd for $C_{19}H_{24}FN_4O_2$ (MH⁺): 359.1883, found: 359.1895.

-49-

Example 22

(7S,8aS)-7-(4-(2-Hydroxyethyl)phenoxy)methyl-2-(5-chloropyridin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




10 A solution of 0.15 g (0.33 mmol) of (7S,8aS)-7-(4-
((methoxycarbonyl)methyl)phenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-
octahydro-pyrrolo[1,2-a]pyrazine (Example 13d) in 15 mL of anhydrous ethyl ether
was added dropwise to an ice-cold suspension of 0.027 g (0.65 mmol) of lithium
aluminum hydride in 15 mL of anhydrous ethyl ether and the mixture stirred for 30 min.
15 The reaction was carefully quenched at 0 °C with 0.027 mL of water, 0.027 mL of
15% sodium hydroxide, and 0.081 mL of water. The precipitate which formed was
filtered through Celite, the filtrate concentrated *in vacuo*, and purification of the
residue by flash silica gel chromatography with 95:5 chloroform:methanol gave 0.13 g
(95%) of the title compound. mp (-HCl) 199-202 °C. HRMS calcd for C₂₁H₂₇ClN₃O₂
20 (MH⁺): 388.1792, found: 388.1807.

Example 23

(7S,8aS)-7-(4-Fluorophenoxy)methyl-2-(6-chloropyrazin-2-yl)-1,2,3,4,6,7,8,8a-
octahydro-pyrrolo[1,2-a]pyrazine

25



30 A mixture of 0.500 g (2.00 mmol) of (7SR,8aSR)-7-(4-fluorophenoxy)methyl-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 3), 1.49 g (10.0 mmol)
of 2,6-dichloropyrazine, and 0.508 g (4.79 mmol) of sodium carbonate in 50 mL of
isoamyl alcohol was heated to reflux for 16 h. The reaction was cooled to ambient


-50-

temperature, the solvent removed *in vacuo*, the residue taken up in ethyl acetate and water, the pH was adjusted to 11 with sodium carbonate and the layers were separated. The aqueous phase was extracted with ethyl acetate, and the combined organic layers were dried (magnesium sulfate), filtered and evaporated. Purification
5 by flash silica gel chromatography with 90:10 ethyl acetate:hexane gave 0.491 g (68%) of the title compound. mp (\cdot HCl) 209-212 $^{\circ}$ C. HRMS calcd for $C_{18}H_{21}ClFN_4O$ (MH $^{+}$): 363.1388, found: 363.1384.

Example 24

(7S,8aS)-7-(4-Fluorophenoxy)methyl-2-(6-chloropyridazin-3-yl)-1,2,3,4,6,7,8,8a-
10 octahydro-pyrrolo[1,2-a]pyrazine

15




A mixture of 0.500 g (2.00 mmol) of (7SR,8aSR)-7-(4-fluorophenoxy)methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 3), 1.49 g (10.0 mmol) of 3,6-dichloropyridazine, and 0.508 g (4.79 mmol) of sodium carbonate in 50 mL of
20 isoamyl alcohol was heated to reflux for 48 h. The reaction was cooled to ambient temperature, the solvent removed *in vacuo*, the residue taken up in ethyl acetate and water, the pH was adjusted to 11 with sodium carbonate and the layers were separated. The aqueous phase was extracted with ethyl acetate, and the combined organic layers were dried (magnesium sulfate), filtered and evaporated. Purification
25 by flash silica gel chromatography with 90:10 ethyl acetate:hexane gave 0.478 g (66%) of the title compound. mp (\cdot HCl) 229 $^{\circ}$ C (dec). HRMS calcd for $C_{18}H_{21}ClFN_4O$ (MH $^{+}$): 363.1388, found: 363.1404.

-51-

Preparation 1

7-Hydroxymethyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




A suspension of 3.20 g (84.3 mmol) of lithium aluminum hydride in 30 mL of dry THF was cooled to 0° C and treated dropwise with a solution of 8.00 g (27.7 mmol) of 7-methoxycarbonyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-1-one (Jones, R. C. F.; Howard, K. J. *J. Chem. Soc., Perkin Trans. 1*, 1993, 2391) in 80 mL of dry THF. After 30 min the reaction was carefully quenched with 3 mL of water, 3 mL of 15% NaOH and 9 of mL water. The mixture was filtered, the filtrate evaporated, the residue taken up in ethyl acetate and washed with brine. The organic layer was dried (magnesium sulfate), filtered, and evaporated to give 6.09 g (89%) of the title compound as a mixture of (7RS,8aSR)- and (7SR,8aSR)- isomers of sufficient purity for use in the next reaction (Preparation 2). m/z (MH⁺) 247.

Preparation 2

(7RS,8aSR)- and (7SR,8aSR)-7-(4-Fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

20



25

A solution of 6.00 g (24.35 mmol) of 7-hydroxymethyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 1), 4.10 g (36.5 mmol) of 4-fluorophenol and 7.70 g (29.4 mmol) of triphenylphosphine in 50 mL dry THF at 0° C was treated dropwise with 4.6 mL (29.3 mmol) of diethyl azodicarboxylate. The reaction was allowed to warm to room temperature and stirred for 24 h. The solvent was evaporated, the residue taken up in ethyl acetate and washed with 1M NaOH (3x). The organic layer was dried (magnesium sulfate), filtered, and evaporated to give the crude product as a dark oil. Purification by flash silica gel chromatography

30

-52-

eluting with 95:5 ethyl acetate:methanol gave 2.27 g (27%) of the (7RS,8aSR)-isomer and 0.410 g (5%) of the (7SR,8aSR)- isomer.


(7RS,8aSR)-7-(4-fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: ^{13}C NMR (CDCl_3): δ 32.0, 35.5, 51.5, 52.5, 56.4, 57.7, 62.7, 62.9, 72.7, 115.45, 115.55, 115.86, 127.0, 128.2, 129.2, 138.3, 155.1, 155.6, 158.8.

(7SR,8aSR)-7-(4-fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine: ^{13}C NMR (CDCl_3): δ 31.6, 35.4, 51.6, 52.5, 57.4, 57.6, 61.7, 62.9, 71.6, 115.40, 115.51, 115.60, 115.91, 127.0, 128.2, 129.2, 138.2, 155.13, 155.15, 155.7, 158.8.

Preparation 3

(7SR,8aSR)-7-(4-Fluorophenoxy)methyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]-pyrazine

15




A mixture of 1.20 g (3.53 mmol) of (7SR,8aSR)-7-(4-fluorophenoxy)methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 2), 30 mL of methanol and 2.5 mL of 5.0 M ammonium formate was treated with an aqueous slurry of 0.15 g of 10% palladium on carbon. The mixture was heated at reflux for 48 h, cooled to ambient temperature, filtered through Celite, and the filtrate was evaporated. The residue was taken up in dilute aqueous ammonium hydroxide and extracted with chloroform (3x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated to give 0.874 g (99%) of the title compound. ^{13}C NMR (CDCl_3): δ 32.0, 34.5, 45.2, 50.9, 53.4, 57.0, 63.7, 72.6, 115.4, 115.5, 115.8, 155.1, 155.6, 158.7. HRMS calcd for $\text{C}_{14}\text{H}_{20}\text{FN}_2\text{O}$ (MH^+): 251.156, found: 251.155.

-53-

Preparation 4

(7RS,8aSR)-7-Hydroxymethyl-7-methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

5




A solution of 3.33 g (11 mmol) of (7RS,8aSR)-7-methoxycarbonyl-7-methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-1-one (Jones, R. C. F.;
10 Howard, K. J. *J. Chem. Soc., Perkin Trans. 1*, 1993, 2391) in 125 mL of dry THF was added dropwise to a stirred suspension of 1.26 g (33 mmol) of lithium aluminum hydride in 125 mL of dry THF. The solution was stirred for 2 h at room temperature, and carefully quenched with 1.26 mL of water, 1.26 mL of 15% NaOH, and 3.78 mL of water. After stirring for 30 min, the mixture was filtered through Celite, dried
15 (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography eluting with 9:1 chloroform:methanol gave 1.67g (58%) of the title compound. ^{13}C NMR (base, CDCl_3): δ 25.3, 39.7, 41.8, 51.1, 51.6, 57.1, 62.3, 62.9, 63.3, 71.3, 127.0, 128.2, 128.7, 129.2, 138.3. m/z (MH^+) 261.

Preparation 5

20 (7RS,8aSR)-7-Hydroxymethyl-7-methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

25



A solution of 1.65 g (6.35 mmol) of (7RS,8aSR)-7-hydroxymethyl-7-methyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 4) in 20 mL of methanol was mixed with 4.44 mL (22.2 mmol) of 5M aqueous ammonium
30 formate, an aqueous slurry of 0.825 g of 10% palladium on carbon was added and the mixture was stirred at ambient temperature overnight. The solution was filtered through Celite and evaporated to give (7RS,8aSR)-7-hydroxymethyl-7-methyl-


-54-

1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine of sufficient purity for use in the next step.

A mixture of (7R,8aSR)-7-hydroxymethyl-7-methyl-1,2,3,4,6, 7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (1.08g, 6.35 mmol), 0.925 g (6.98 mmol) of 2-chloro-5-fluoropyrimidine (Dunaïskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), and 1.48 g (13.96 mmol) of sodium carbonate and 65 mL of water was heated at 90 °C for 72 h. The solution was cooled and extracted with chloroform (3x). The combined organic layers were dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography eluting with chloroform:methanol (95:5) gave 0.80 g (47%) of the title compound. ¹³C NMR (CDCl₃): δ 25.5, 39.5, 41.7, 43.3, 48.7, 51.2, 62.3, 63.8, 71.0, 144.9, 145.2, 149.8, 153.1, 158.9. m/z (MH⁺) 267.

Preparation 6

(7R,8aS)-7-Hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



Step A

A solution of 9.75 g (42.0 mmol) of (7R,8aS)-7-hydroxy-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Diafi, L.; et al. *J. Het. Chem.*, 1990, 27, 2181) and 29.4 mL of 5M ammonium formate in 140 mL of methanol was treated with an aqueous slurry of 4.9 g of 10% palladium on carbon and the mixture was stirred at ambient temperature for 18 h. The mixture was filtered through Celite and evaporated to give (7R,8aS)-7-hydroxy-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine as a clear oil.

Step B

A mixture of the crude (7R,8aS)-7-hydroxy-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine, 9.96 g (52.5 mmol) of 2-chloro-5-fluoropyrimidine (Dunaïskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), 13.4 g (126 mmol) of sodium carbonate and 450 mL of water was heated at 95 °C for 72 h. The mixture was cooled, extracted with chloroform (2x), and the combined organic phase was dried

-55-


(magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 95:5 ethyl acetate:methanol gave 8.54 g (85%) of the title compound. ^{13}C NMR (base, CDCl_3): δ 39.1, 43.7, 48.7, 51.0, 60.2, 62.9, 69.4, 144.95, 145.24, 149.9, 153.2, 158.9. m/z (MH^+) 239.

5

Preparation 7

(7R,8aS)-7-Methanesulfonyloxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

10




A solution of 1.00 g (4.20 mmol) of (7R,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 6) and 0.644 mL (4.62 mmol) of triethylamine in 40 mL of methylene chloride was chilled to 0 °C and 0.34 mL (4.41 mmol) of methanesulfonyl chloride in 20 mL of methylene chloride was added slowly. After 30 min, water was added and the pH adjusted to 10 with 1M NaOH. The layers were separated, the organic phase was washed with water (2x), dried (magnesium sulfate), filtered and evaporated to give 1.15 g (86%) of the title compound of sufficient purity for use in subsequent reactions. m/z (MH^+) 317.

20

Preparation 8

7-Methoxycarbonyl-2-phenylmethyl-1,2,3,4,8,8a-hexahydro-pyrrolo[1,2-a]pyrazin-1-one

25



A solution of 8.6 g (27 mmol) of dimethyl 2-phenylmethyl-2,3,5,6,7,7a-hexahydro-1H-pyrrolo[1,2-c]imidazol-5,7-dioate (Jones, R. C. F.; Howard, K. J. J. *Chem. Soc. Perkin Trans. 1*, 1993, 2391) and 2.0 mL (13 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) in 150 mL of methanol was heated at reflux for 16 h. The solvent was evaporated, the residue was dissolved in ethyl acetate and

30

-56-


washed with water (2x), dried (magnesium sulfate), filtered and evaporated to give 6.45 g (84%) of the title compound. ^{13}C NMR (CDCl_3): δ 32.9, 43.9, 45.6, 50.5, 50.8, 63.1, 106.0, 127.7, 128.0, 128.8, 136.2, 148.1, 165.7, 170.0. HRMS calcd for $\text{C}_{16}\text{H}_{19}\text{N}_2\text{O}_3$ (MH^+): 287.1396, found: 287.1406.

5

Preparation 9

(7SR,8aSR)-7-Methoxycarbonyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-1-one

10



15


A mixture of 40.0 g (140 mmol) of 7-methoxycarbonyl-2-phenylmethyl-1,2,3,4,8,8a-hexahydro-pyrrolo[1,2-a]pyrazin-1-one (Preparation 8) and 10 g of 5% Pd on charcoal in 600 mL of ethyl acetate was shaken in a Parr apparatus under 50 psi of hydrogen gas for 6.5 h. The mixture was filtered through Celite, and the filtrate was evaporated to give 38.4 g (95%) of the title compound. ^{13}C NMR (CDCl_3): δ 31.8, 41.4, 44.8, 46.5, 49.6, 52.1, 55.2, 64.0, 127.5, 128.1, 128.7, 128.8, 136.6, 169.6, 174.8. HRMS calcd for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}_3$ (MH^+): 289.1552, found: 289.1549.

20

Preparation 10

(7SR,8aSR)-7-Hydroxymethyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

25



30

A three-neck flask fitted with a reflux condenser and dropping funnel was charged with 100 mL of dry THF and 2.4 g (63 mmol) of lithium aluminum hydride (LAH), and a solution of 6.0 g (21 mmol) of (7SR,8aSR)-7-methoxycarbonyl-2-phenylmethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-1-one (Preparation 9) in 60 mL of dry THF was placed in the dropping funnel. The LAH suspension was heated to reflux and the ester solution was added over a 30-60 min period. Reflux


-57-

continued for another 4 h, the reaction was cooled in an ice bath and quenched by careful addition of 2.4 mL of water, 2.4 mL of 15% sodium hydroxide, and 7.2 mL of water. Stirring continued until a white precipitate formed, the mixture was filtered through Celite, and the filtrate was evaporated to give 5.0 g (97%) of the title

5 compound of sufficient purity for use in subsequent reactions. ^{13}C NMR (CDCl_3): δ 31.1, 37.2, 51.3, 52.6, 57.4, 62.8, 62.9, 67.4, 127.0, 128.2, 129.2, 138.2. HRMS calcd for $\text{C}_{15}\text{H}_{23}\text{N}_2\text{O}$ (MH^+) 247.1810, found: 247.1800.

Preparation 11

(7S,8aS)-7-Methoxycarbonyl-1,2,3,4,6,7,8,8a-octahydro-
10 pyrrolo[1,2-a]pyrazin-1-one



15

A mixture of 77.3 g (16 mmol) of N-benzyloxycarbonyl-*cis*-4-carboxy-L-proline dimethyl ester (Bridges, R. J; et al. *J. Med. Chem.*, 1991, 34, 717) and 169 mL (55 mmol) of 5M ammonium formate in 1000 mL of methanol was treated with an aqueous slurry of 15.5 g of 10% palladium on carbon. After 4 h, the mixture was
20 filtered through Celite, the filtrate was concentrated to about 500 mL, saturated with sodium bicarbonate, and extracted with chloroform (5x). The aqueous layer was saturated with sodium chloride and extracted with chloroform (3x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated to give 39.8 g (88%) of *cis*-4-carboxy-L-proline dimethyl ester.

25 A mixture of 39.8 g (213 mmol) of *cis*-4-carboxy-L-proline dimethyl ester, 40.2 g (213 mmol) of 2-(phthalimido)acetaldehyde (Preparation 12), and 17.5 g (213 mmol) of anhydrous sodium acetate in 2150 mL of dry methylene chloride was treated with 67.7 g (319 mmol) of sodium (triacetoxy)borohydride in small portions over a 1 h period. The solution was stirred for 16 h, water was added, and the pH
30 adjusted to 10 with 1M sodium hydroxide. The organic phase was separated, dried (magnesium sulfate), filtered and evaporated to give 73.8 g (96%) of N-(2-(phthalimido)ethyl)-*cis*-4-carboxy-L-proline dimethyl ester.

-58-


A solution of 73.8 g (205 mmol) of N-(2-(phthalimido)ethyl-*cis*-4-carboxy-L-proline dimethyl ester and 44.1 mL (513 mmol) of 40% aqueous methyl amine in 3100 mL of methanol was stirred at ambient temperature for 16 h. The solvent was evaporated and the crude product purified by flash silica gel chromatography starting
5 with 4:1 diethyl ether:methanol and ending with 2:1 diethyl ether: methanol to give 37.0 g (91%) of the title compound of sufficient purity for use in subsequent reactions. *m/z* (MH⁺) 199.

Another sample of the title compound was purified further by flash silica gel chromatography with ethyl ether:methanol (9:1 to 8:2) and formed a white solid on
10 standing. mp 70-74 °C. ¹³C NMR (d₆-DMSO): δ 32.2, 40.5, 42.4, 46.6, 52.6, 55.8, 64.0, 173.9, 176.2. HRMS calcd for C₉H₁₅N₂O₃ (MH⁺):199.1083, found: 199.1091.

Preparation 12

2-(Phthalimido)acetaldehyde

15




A solution of 50.0 g (190 mmol) of 2-(phthalimido)acetaldehyde diethyl acetal
20 in 300 mL of toluene was treated with 150 mL of 50% aqueous trifluoroacetic acid and the mixture was stirred vigorously at ambient temperature for 72 h. The solution was concentrated *in vacuo* and 100 mL of ethyl acetate was added. The white precipitate was filtered off and washed with ice-cold ethyl acetate (100 mL) to give 30.0 g (78%) of the title compound.

25

Preparation 13

(7*S*,8*aS*)-7-Hydroxymethyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8*a*-octahydro-pyrrolo[1,2-*a*]pyrazine

30



-59-

Step A

A solution of 1.75 g (8.84 mmol) of (7S,8aS)-7-methoxycarbonyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-1-one (Preparation 11) in 100 mL of THF was added dropwise to a suspension of 0.67 g (18 mmol) of lithium aluminum hydride in 100 mL of refluxing THF. After stirring for 1 h, the solution was cooled and carefully quenched with 0.67 mL of water, 0.67 mL of 15% sodium hydroxide, and 2.0 mL of water. The precipitate was filtered and the filtrate was concentrated to give 1.4 g of (7S,8aS)-7-hydroxymethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine of sufficient purity for use in subsequent reactions.

Step B


A mixture of 1.38 g (8.84 mmol) of (7S,8aS)-7-hydroxymethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine, 1.30 g (9.87 mmol) of 2-chloro-5-fluoropyrimidine (Dunaïskis, A.; et al. *Org. Prep. Proc. Int.*, 1995, 27, 600-602), and 2.85 g (26.9 mmol) of sodium carbonate and 90 mL of water was heated at 95 °C for 16 h. The solution was cooled and extracted with chloroform (2x), the combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 9:1 chloroform:methanol gave 0.97 g (43%) of the title compound. mp 123-124 °C. m/z (MH⁺) 253. ¹³C NMR (CDCl₃): δ 30.9, 37.1, 43.8, 48.8, 51.3, 57.5, 62.6, 67.2, 144.9, 145.2, 149.9, 153.2, 158.8.

20

Preparation 14

(7S,8aS)-7-Hydroxymethyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine

25



30


A mixture of 0.50 g (3.2 mmol) of (7S,8aS)-7-hydroxymethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 13, Step A), 2.37 g (16.0 mmol) of 2,5-dichloropyridine, 0.85 g (8.0 mmol) of sodium carbonate and 35 mL of isoamyl alcohol was refluxed for 48 h. The hot solution was filtered and the filtrate evaporated. Purification by flash silica gel chromatography with 9:1 chloroform:methanol gave 0.25 g (29%) of the title compound. ¹³C NMR (CDCl₃): δ

-60-

31.1, 37.3, 44.7, 49.7, 51.2, 57.2, 62.5, 66.9, 107.9, 119.9, 137.1, 146.1, 157.6. m/z (MH⁺) 268.

Preparation 15


(7S,8aS)-7-Hydroxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



A solution of 1.58 g (4.61 mmol) of (7S,8aS)-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate (Example 9) in 200 mL of methanol was treated with 50 mL of 15% aqueous sodium hydroxide. After 30 min, the solvent was removed and the residue partitioned between water and ethyl acetate. The layers were separated, the organic phase was dried (magnesium sulfate), filtered and evaporated to give 0.922 g (84%) of the title compound. ¹³C NMR (CDCl₃): δ 39.4, 43.7, 49.0, 51.2, 62.2, 63.6, 69.9, 144.9, 145.2, 149.9, 153.2, 158.9. m/z (MH⁺) 239.

Preparation 16

(7S,8aS)-7-Hydroxymethyl-2-(4-cyanophenyl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine




A mixture of 1.5 g (9.6 mmol) of (7S,8aS)-7-hydroxymethyl-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 13, Step A), 1.75 g (14.4 mmol) of 4-fluorobenzonitrile and 2.04 g (19.2 mmol) of sodium carbonate in 10 mL of DMSO was heated at 80 °C for 16 h. The solution was cooled to room temperature, diluted with water and extracted with 1:1 ethyl acetate:diethyl ether (3x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 9:1 chloroform: methanol gave 0.925 g (37%) of

-61-

the title compound. ^{13}C NMR (CDCl_3): δ 31.1, 37.2, 46.7, 51.0, 51.8, 57.6, 62.3, 67.3, 100.1, 114.3, 120.0, 133.5, 153.4. m/z (MH^+) 258.

Preparation 17


(7R,8aS)-7-Hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine



A mixture of 3.81 g (26.8 mmol) of (7R,8aS)-7-hydroxy-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine (Preparation 6, Step A), 19.8 g (134 mmol) of 2,5-dichloropyridine, 7.10 g (67.0 mmol) of sodium carbonate and 275 mL of isoamyl alcohol was heated at reflux for 72 h. The mixture was cooled to about 100 °C, filtered hot, and the filtrate concentrated *in vacuo*. Purification of the residue by flash silica gel chromatography with 9:1 chloroform:methanol gave 0.63 g (9%) of the title compound of sufficient purity for use in subsequent reactions. m/z (MH^+) 254.

Preparation 18

(7S,8aS)-7-Hydroxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine




A solution of 0.52 g (1.45 mmol) of (7S,8aS)-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate (Example 17) in 50 mL of methanol was treated with 50 mL of 15% aqueous sodium hydroxide and stirred at ambient temperature for 30 min. The solvent was concentrated by half *in vacuo* and the residue extracted with chloroform (3x). The combined organic phase was dried (magnesium sulfate), filtered and evaporated. Purification by flash silica gel chromatography with 9:1 chloroform:methanol gave 0.25 g (68%) of the title compound. mp (base) 167-168 °C. m/z (MH^+) 254.

-62-

Claims

1. A compound of formula I



wherein

R_1 is phenyl, naphthyl, benzoxazolonyl, indolyl, indolonyl, benzimidazolyl, quinolyl, furyl, benzofuryl, thienyl, benzothienyl, oxazolyl, benzoxazolyl;

10 R_2 is H or (C₁-C₆)alkyl;


R_3 is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, or pyridazinyl;

R_4 is H or (C₁-C₆)alkyl;

R_5 is H or (C₁-C₆)alkyl;

wherein each group of R_1 and R_3 may be independently and optionally
 15 substituted with one to four substituents independently selected from the groups consisting of fluoro, chloro, bromo, iodo, cyano, nitro, thiocyno, -SR₄, -SOR₄, -SO₂R₄, -NHSO₂R₄, -(C₁-C₆)alkoxy, -NR₄R₅, -NR₄COR₅, -CONR₄R₅, phenyl, -COR₄, -COOR₄, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl substituted with one to six halogens, -(C₃-C₆)cycloalkyl, and trifluoromethoxy;

20 X is O, S, SO, SO₂, NR₄, C=O, CH(OH), CHR₄,



m is 0, 1 or 2;

n is 0, 1 or 2;

all stereoisomers thereof; or

25 a pharmaceutically acceptable salt thereof.

2. A compound of claim 1 wherein

R_1 is phenyl, naphthyl, benzoxazolonyl, indolyl, indolonyl, benzimidazolyl, or quinolyl;

wherein R_1 and R_3 may be independently substituted with up to three
 30 substituents independently selected from the group consisting of fluoro, chloro, bromo, iodo, cyano, -NR₄R₅, -(C₁-C₆)alkoxy, -COOR₄, -CONR₄R₅, -(C₁-C₆)alkyl, -

-63-

(C₁-C₆)alkyl substituted with one to six halogens, -(C₃-C₆)cycloalkyl, and trifluoromethoxy;

R₂ is H or CH₃;

X is O, C=O, CHOH, -C(=O)O-, or CH₂;

5 m is 0 or 1;

n is 0 or 1; or

a pharmaceutically acceptable salt thereof.

3. A compound of claim 2 wherein

R₁ is phenyl or substituted phenyl;

10 R₃ is substituted or unsubstituted phenyl, pyridinyl, or pyrimidinyl;

X is O, -C(=O)O-, or CH₂; or

a pharmaceutically acceptable salt thereof.

4. A compound of claim 3 wherein

R₂ is H;

15 X is O;

m is 0;

n is 1; or

a pharmaceutically acceptable salt thereof.

5. A compound of claim 3 wherein

20 R₂ is H;

X is O;

m is 1;

n is 0; or

a pharmaceutically acceptable salt thereof.

25 6. A compound of claim 3 wherein

R₂ is H;

X is -C(=O)O-;

m is 0;

n is 0; or

30 a pharmaceutically acceptable salt thereof.

7. A compound of claim 4 wherein

R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;

-64-

- R₃ is chloropyridinyl; or
a pharmaceutically acceptable salt thereof.
8. A compound of claim 4 wherein
R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;
5 R₃ is fluoropyrimidinyl; or
a pharmaceutically acceptable salt thereof.
9. A compound of claim 5 wherein
R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;
R₃ is chloropyridinyl; or
10 a pharmaceutically acceptable salt thereof.
10. A compound of claim 5 wherein
R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;
R₃ is fluoropyrimidinyl; or
a pharmaceutically acceptable salt thereof.
- 15 11. A compound of claim 6 wherein
R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;
R₃ is chloropyridinyl; or
a pharmaceutically acceptable salt thereof.
12. A compound of claim 6 wherein
20 R₁ is fluorophenyl, difluorophenyl, or cyanophenyl;
R₃ is fluoropyrimidinyl; or
a pharmaceutically acceptable salt thereof.
13. A compound of claim 7 wherein
R₃ is 5-chloro-pyridin-2-yl; or
25 a pharmaceutically acceptable salt thereof.
14. A compound of claim 8 wherein
R₃ is 5-fluoro-pyrimidin-2-yl; or
a pharmaceutically acceptable salt thereof.
15. A compound of claim 9 wherein
30 R₃ is 5-chloro-pyridin-2-yl; or
a pharmaceutically acceptable salt thereof.
16. A compound of claim 10 wherein

-65-

R₃ is 5-fluoro-pyrimidin-2-yl-; or
a pharmaceutically acceptable salt thereof.

17. A compound of claim 11 wherein

R₃ is 5-chloro-pyridin-2-yl-; or

5 a pharmaceutically acceptable salt thereof.

18. A compound of claim 12 wherein

R₃ is 5-fluoro-pyrimidin-2-yl-; or

a pharmaceutically acceptable salt thereof.

19. A compound of claim 1 which is (7S,8aS)-7-(4-fluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

20. A compound of claim 1 which is (7S,8aS)-7-(3,5-difluorophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

21. A compound of claim 1 which is (7S,8aS)-7-(3-cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

15 22. A compound of claim 1 which is (7S,8aS)-7-(4-cyanophenoxy)methyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

23. A compound of claim 1 which is (7S,8aS)-7-(4-fluorobenzyl)oxy-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

20 24. A compound of claim 1 which is (7S,8aS)-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate.

25. A compound of claim 1 which is (7S,8aS)-7-(4-fluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

26. A compound of claim 1 which is (7S,8aS)-7-(3,5-difluorophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

25 27. A compound of claim 1 which is (7S,8aS)-7-(3-cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

28. A compound of claim 1 which is (7S,8aS)-7-(4-cyanophenoxy)methyl-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

29. A compound of claim 1 which is (7S,8aS)-7-(4-fluorobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

30. A compound of claim 1 which is (7S,8aS)-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazin-7-yl benzoate.

-66-

31. A compound of claim 1 which is (7S,8aS)-7-(3-cyanobenzyl)oxy-2-(5-fluoropyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

32. A method for treating or preventing disorders of the dopamine system in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

33. A method for treating or preventing psychotic disorders such as affective psychosis, schizophrenia, and schizoaffective disorders in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

34. A method for treating or preventing movement disorders such as extrapyramidal side effects from neuroleptic agents, neuroleptic malignant syndrome, tardive dyskinesia, or Gilles De La Tourette's syndrome in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

35. A method for treating or preventing movement disorders such as Parkinson's disease or Huntington's disease in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

36. A method for treating or preventing gastrointestinal disorders such as gastric acid secretion in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

37. A method for treating or preventing gastrointestinal disorders such as emesis in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

38. A method for treating or preventing chemical abuse, chemical dependencies or substance abuse in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

-67-

39. A method for treating or preventing vascular and cardiovascular disorders such as congestive heart failure and hypertension in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
40. A method for treating or preventing ocular disorders in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
41. A method for treating or preventing sleep disorders in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
42. A pharmaceutical composition for treating or preventing disorders of the dopamine system in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
43. A pharmaceutical composition for treating or preventing psychotic disorders such as affective psychosis, schizophrenia, and schizoaffective disorders in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
44. A pharmaceutical composition for treating or preventing movement disorders such as extrapyramidal side effects from neuroleptic agents, neuroleptic malignant syndrome, tardive dyskinesia, or Gilles De La Tourette's syndrome in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.
45. A pharmaceutical composition for treating or preventing movement disorders such as Parkinson's disease or Huntington's disease in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

-68-

46. A pharmaceutical composition for treating or preventing gastrointestinal disorders such as gastric acid secretion in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

5 47. A pharmaceutical composition for treating or preventing gastrointestinal disorders such as emesis in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.


48. A pharmaceutical composition for treating or preventing chemical abuse,
10 chemical dependencies or substance abuse in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

49. A pharmaceutical composition for treating or preventing vascular and
15 cardiovascular disorders such as congestive heart failure and hypertension in a mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

50. A pharmaceutical composition for treating or preventing ocular disorders in a
20 mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

51. A pharmaceutical composition for treating or preventing sleep disorders in a
25 mammal, comprising administering to said mammal an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder.

52. A compound of formula IV



wherein

n is 0 or 1;

-69-

Y is CH or N;

Z is chloro or fluoro;

all stereoisomers thereof;

which are useful intermediates for the preparation of compounds of formula I.

- 5 53. A compound of claim 52 which is (7S,8aS)-7-hydroxymethyl-2-(5-chloropyridin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.
54. A compound of claim 52 which is (7S,8aS)-7-hydroxymethyl-2-(5-fluoro-pyrimidin-2-yl)-1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.
55. A compound of claim 52 which is (7R,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-
10 1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.
56. A compound of claim 52 which is (7R,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.
57. A compound of claim 52 which is (7S,8aS)-7-hydroxy-2-(5-chloropyridin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.
- 15 58. A compound of claim 52 which is (7S,8aS)-7-hydroxy-2-(5-fluoropyrimidin-2-yl)-
1,2,3,4,6,7,8,8a-octahydro-pyrrolo[1,2-a]pyrazine.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 96/01192

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D487/04 A61K31/495 //(C07D487/04,241:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 380 217 A (PFIZER) 1 August 1990 see page 1, line 22 - line 23; claim 1 ---	1,32
P,A	WO 96 10571 A (PFIZER) 11 April 1996 see claims 1,16 -----	1,32

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

30 January 1997

Date of mailing of the international search report

1 0. 02. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 96/ 01192

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 32-41 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/IB 96/01192

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-380217	01-08-90	WO-A- 9008148	26-07-90
		AP-A- 128	20-03-91
		BG-A- 51046	15-01-93
		CA-A,C 2008108	23-07-90
		CN-A,B 1044464	08-08-90
		EG-A- 19306	31-01-95
		ES-T- 2053094	16-07-94
		FI-B- 95381	13-10-95
		IE-B- 62670	08-03-95
		IL-A- 93073	30-05-94
		JP-A- 2233681	17-09-90
		JP-B- 6043427	08-06-94
		NO-B- 175978	03-10-94
		PL-B- 163836	31-05-94
		PT-B- 92919	29-12-95
		RU-C- 2015977	15-07-94
		WO-A- 9008144	26-07-90
		US-A- 5122525	16-06-92

WO-A-9610571	11-04-96	AU-A- 3189395	26-04-96
		BR-A- 9504230	06-08-96
